

Wassie K. B. B.

BIOLOGICAL JOURNAL of the Linnean Society

VOLUME 58

NUMBERS 1-4

PAGES 1-500

THE LINNEAN SOCIETY OF LONDON
BURLINGTON HOUSE, PICCADILLY, LONDON W1V 0LQ
1996

Dates of publication of the numbers included in Vol. 58:

No. 1, pp. 1–123, 15 May 1996

No. 2, pp. 125–246, 24 June 1996

No. 3, pp. 247–369, 12 July 1996

No. 4, pp. 371–500, 12 September 1996

CONTENTS

Pages

ANDERSSON, S., Flower-fruit size allometry at three taxonomic levels in <i>Crepis</i> (Asteraceae)	401-407
BERGLUND, A., BISAZZA, A. and PILASTRO, A., Armaments and ornaments: an evolutionary explanation of traits of dual utility	385-399
BJORKLUND, M., Similarity of growth among Great tits (<i>Parus major</i>) and Blue tits (<i>P. caeruleus</i>)	343-355
COOK, I. M. and GRAHAM, C. S., Evenness and species number in some moth populations	75-84
COSTA, J. T., McDONALD, J. H. and PIERCE, N. E., The effect of ant association on the population genetics of the Australian butterfly <i>Jalmenus evagoras</i> (Lepidoptera: Lycaenidae)	287-306
DUFFY, J. E., Species boundaries, specialization, and the radiation of sponge-dwelling alpheid shrimp	307-323
FRYER, G., Reflections on arthropod evolution	1-55
GREENE, E., Effect of light quality and larval diet on morph induction in the polymorphic caterpillar <i>Nemoria arizonaria</i> (Lepidoptera: Geometridae)	277-285
GUINARD, R., Use of a multivariate model using allele frequency distributions to analyse patterns of genetic differentiation among populations	173-195
HARLIN, M., Biogeographic patterns and the evolution of eurentic nemertean	325-342
HEWITT, G. M., Some genetic consequences of ice ages, and their role in divergence and speciation	247-276
JACKSON, K., BUTLER, D. G. and BROOKS, D. R., Habitat and phylogeny influence salinity discrimination in crocodilians: implications for osmoregulatory physiology and historical biogeography	371-383
JOHNSON, M. S. and BLACK, R., Geographic cohesiveness versus associations with habitat: genetic subdivision of <i>Bembicium vittatum</i> Philippi (Gastropoda: Littorinidae) in the Houtman Abrolhos Islands	57-74
JOHNSTONE, R. A. and HURST, G. D. D., Maternally inherited male-killing micro-organisms may confound interpretation of mitochondrial DNA variability	453-470
LARSEN, T., SORDAHL, T. A., and BYRKJEDAL, I., Factors related to aggressive nest protection behaviour: a comparative study of Holarctic waders	409-439
MICHELSSEN, V., First reliable record of a fossil species of Anthomyiidae (Diptera), with comments on the definition of recent and fossil clades in phylogenetic classification	441-451
PARSONS, P. A., Conservation strategies: adaptation to stress and the preservation of genetic diversity	471-482
PRITCHARD, G., HARDER, L. D. and MUTCH, R. A., Development of aquatic insect eggs in relation to temperature and strategies for dealing with different thermal environments	221-244
ROBINSON, J. V. and ALLGEYER, R., Covariation in life-history traits, demographics and behaviour in ischnuran damselflies: the evolution of monandry	85-98
RUEDI, M., Phylogenetic evolution and biogeography of Southeast Asian shrews (genus <i>Crocodyra</i> : Soricidae)	197-219
SNEATH, P. H. A., The statistical reliability of synapomorphic analyses	483-497
STEWART, C. N., Jr., ROSSON, G., SHIRLEY, B. W. and PORTER, D. M., Population genetic variation in rare and endangered <i>Ilamnia</i> (Malvaceae) in Virginia	357-369
TRIGO, J. R., BROWN, K. S., Jr., WITTE, L., HARTMANN, T., ERNST, L. and SOARES BARATA, L. E., Pyrrolizidine alkaloids: different acquisition and use patterns in Apocynaceae and Solanaceae feeding ithomiine butterflies (Lepidoptera: Nymphalidae)	99-123
WESTMORELAND, D. and KILTIE, R. A., Egg crypsis and clutch survival in three species of blackbirds (Icteridae)	159-172
WILLIAMS, P. H., PRANCE, G. T., HUMPHRIES, C. J., and EDWARDS, K. S., Promise and problems in applying quantitative complementary areas for representing the diversity of some Neotropical plants (families Dichapetalaceae, Lecythidaceae, Caryocaraceae, Chrysobalanaceae and Proteaceae)	125-157

BOOK REVIEWS

<i>Darwinism Evolving. Systematics Dynamics and the Genealogy of Natural Selection</i> , by D. J. Depew and B. H. Weber. (M. F. CLARIDGE)	499-500
<i>Ecological versatility and Community Ecology</i> , by Ralph C. Mac Nally (M. O. HILL)	245
<i>Exploring Evolutionary Biology</i> , by M. Slatkin. Readings from <i>American Scientist</i> . (R. J. BERRY)	500
<i>Functional Morphology in Vertebrate Palaeontology</i> , ed. J. J. Thomason (P. FOREY)	245-246
<i>Speciation and the Recognition Concept, Theory and Application</i> , by D. M. Lambert and H. G. Spencer. (M. F. CLARIDGE)	499

INDEX

Compiled by Stanley Thorley

Abbreviations:

App.—Appendix

Fig.—Figure

n.—note

Tab.—Table

*—signifies a keyword

- Aat (E.C. 2.6.1.1)
butterfly-ant mutualism
 Aat-1 291, 295 Tabs 2–3
 Aat-1 and *Aat-1* 299 Tab. 5
sponge-dwelling alpheid shrimps 310, 312–314 &
 Fig. 1 & Tab. 1
- Acacia* (Leguminosae)
host plants: butterfly-ant mutualism 290, 292 Tab.
 1, 300–301, 303
 A. dealbata 292 Tab. 2
 A. decurrens 291–292 & Tab. 2
 A. filicifolia 291–292 & Tab. 1
 A. harpophylla (brigalow) 288
 A. melanoxylon 291–292 & Tab. 2
pyrrolizidine alkaloids in ithomiine butterflies: *A.*
 plumosa 108 Tab. 3, 110 Tab. 5
- Acanthaceae: pyrrolizidine alkaloids in ithomiine
 butterflies 106–109 Figs 2–3 & Tabs 1 & 4
 see also *Dicliptera* and *Ruellia*
- acanthuroid fishes: cladistic biogeography of the
 Eureptantia 325–326, 330, 332–333, 336–337
 Figs 5–6 & Tab. 3, 340
- ACCTRAN: cladistic biogeography of the Eureptantia
 339
- Acheareana wau* (social spiders): acoustic signals 391–392
 Tab. 1
- Acinonyx jubatus* (cheetahs): genetic stress 472–473
- Acioa*: complementary areas for diversity
 A. edulis 151 App. 2
 A. guianensis 147 App. 2
 A. schultesii 147 App. 2
 A. somnolens 147 App. 2
- Acipenser transmontanus* (white sturgeon): molecular data
 272 App.
- Acch-1* (E.C. 4.2.1.3.): butterfly-ant mutualism 291,
 295 Tabs 2–3
- Acch-1* 299 Tab. 5
- Acraeinae: pyrrolizidine alkaloids in ithomiine
 butterflies 109, 119
- Acracini 120
- Actinote* 109
- Acrocephalus arundinaceus* (great reed warblers): acoustic
 signals 391–393 & Tab. 1
- Acrothoracica: eyeless adults 42
- Actinia tenebrosa* (anemones): intertidal communities 61,
 64 Tab. 1
- Actitis*: aggressive nest defence
 A. hypoleucos 438 App.
 A. macularia 438 App.
- ADA (EC 3.5.4.4.): Southeast Asian shrews 200, 203
 Ada 204–206 Tabs 1–3
- Adalia bipunctata* (two-spot ladybirds): cytoplasmic male-
 killers 459, 462–463, 467–468
- adaptation: development of aquatic insect eggs
 221–225 & Fig. 1, 227–229 7 Fig. 4, 232–233
- *adaptation: genetics of ice ages 247–248, 262–266 &
 Fig. 7
- *adaptive coloration: *Nemoria arizonaria* (emerald
 moths) 277
- ADE (Analyse des Donnée Environnementales) software:
 multivariate genetic differentiation 192
- adenosine deaminase locus: Soay sheep 474
- Adenostemma swartzii* (Asteraceae): ithomiine visits 108
 Tab. 3, 110 Tab. 5
- ADH (EC 1.1.1.1.): Southeast Asian shrews 200
 Adh 205 Tab. 2
- Aedes*
 Culicidae: development of aquatic insect eggs 228
 A. aegypti 228, 231, 242 App. 2
 A. sticticus 228, 23, 242 App. 2
 A. vexans 242 App. 2
 molecular analysis 46
- Acquifurcata: division of the eureptantian nemerteans
 326, 338
- **Aeria olena* (ithomiine butterflies): pyrrolizidine
 alkaloids 99, 101, 104–108 & Tabs 1–3, 111,
 115, 118–120
- Aeshna* (Odonata): development of aquatic insect eggs
 233
- Africa
 genetics of ice ages 257, 264, 275 App.
 North 249, 251, 257
 Southeast Asian shrews 197, 199, 203, 213
- Agelaius phoeniceus* (red-winged blackbirds)
 acoustic and visual signals 391–393 & Tab. 1

- egg crypsis 159–162 & Fig. 1, 164–171 & Tabs 1–3 & Figs 2–4
- Agelas clathrodes* (sponges): Belize 310, 314, 316, 318 Fig. 4
- Ageratum conyzoides* (Eupatoriaceae): Asteraceae): pyrrolizidine alkaloids 106–108 Fig. 2 & Tabs 1 & 3, 113 & Tab. 9, 117
- *aggressive nest defence: protection by waders 409–414, 418–429 & Figs 3–6 & Tabs 1–2, 436–439 App
- Aglaspidida: Cambrian arthropod 32
- AK (EC 2.7.4.3.)
butterfly-ant mutualism
Ak 291, 297, 295 Tabs 2–3
Ak 299 Tab. 5
Southeast Asian shrews
Ak-1 204–205 Tabs 1–2
AK-1 and AK-2 200
AK-2 203
- Akodontine tribe (murid rodents): molecular data 274 App.
- ALB: Southeast Asian shrews 200, 203
Alb 204–206 Tabs 1–3
- Albrolhos archipelago (Australia): *Bembicium vittatum* (snails) 57–59, 64–66 & Fig. 3, 69–70
- alder pollen cores: Europe 250
- alkaloids, pyrrolizidine (PAs) *see* pyrrolizidine alkaloids in ithomiine butterflies
- alleles
butterfly-ant mutualism: *Jalmenus evagoras* (Lycaenidae) 291, 293–295 & Tabs 3 & 5, 296–299 & Fig. 5
genetics of ice ages 253–255 & Fig. 2, 260
geographic cohesiveness in an intertidal snail 57–58, 62, 65–68 & Fig. 3 & Tab. 2, 70–71
multivariate genetic differentiation 173–177 & Fig. 1, 179–185 & Figs 1 & 4, 188–192
Southeast Asian shrews 201, 203–207 & Tabs 1–3, 213, 215
specialization and speciation in shrimps 308–310, 312–314 & Fig. 1 & Tab. 1, 319–320
- *allometry: growth similarities in tits 343–352 & Tab. 1–5 & Fig. 1
- allometry, flower-fruit size: *Crepis* (Asteraceae) 401–407 & Figs 12
- *allozymes: geographic cohesiveness in an intertidal snail 57–59, 65–70 & Figs 3–5 & Tabs 2–3, 72
- *allozymes: SE Asian shrews 197, 201, 203, 214–215
- allozymes
genetics of ice ages 257–258, 260, 264
Nucella lapillus (dogwhelks): multivariate genetic differentiation 180–181, 187
see also Est-3, Lap-1, Lap-2, Mdh-1, Pep-1, Pep-2, Pgm-1 & Pgm-2
specialization and speciation in shrimps 307–309, 316 Fig. 2, 319–321
- Alpheidae: sponge-dwelling shrimps 307–308
Alpheus 314
- Alps: genetics of ice ages 249, 251–253, 258–259 & Fig. 5, 262, 264–265 & Fig. 7
- Althernanthera ficoidea* (Amaranthaceae): *Aeria olen*a visits 106–107 Fig. 2 & Tab. 1
- amabiline (pyrrolizidine alkaloid): ithomiine butterflies 99, 102 Fig. 1, 113–114
- Amaranthaceae: *Aeria olen*a visits 106–107 Fig. 2 & Tab. 1
- Althernanthera ficoidea* 106–107 Fig. 2 & Tab. 1
- Amazonia: complementary areas for diversity 125, 127, 134, 138
- amber, Dominican: first fossil anthomyid 441–443 & Fig. 1
- Ambon (Indonesian archipelago): Eureptantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
- America *see* North and South America
- Amia calva* (bowfins): molecular data 272 App.
- Anmodramus maritimus* ('seaside sparrows'): molecular data 272 App.
- Anmodramus* (sparrows): ornaments are armaments
A. maritimus (Scott's seaside) 391–392 Tab. 1
A. swannarum (grasshopper) 393
- *AMOVA (analysis of molecular variance): RAPD variation in *Iliamna* 357, 359–361, 364–365 & Fig. 2
- Amphibia
complex colonization scheme: *Bufo marinus* 188
see also frogs and toads
Amphinenura (Nemouridae): development of aquatic insect eggs
A. banksi 232, 238 App. 2
A. standfussi 238 App. 2
A. sulcipectus 238 App. 2
- Amphipoda: eye structure 28
- Amphicusta maya* (gregarious crickets): acoustic signals 391–392 Tab. 1
- analyses
aggressive nest protection by waders 410, 413–418, 423–429, 436–439 App.
see also chi-squares, Mann-Whitney U-tests, regressions and Spearman rank correlation tests
butterfly-ant mutualism 290–295 & Fig. 2 & Tabs 2–3
see also bootstrap resampling, jackknife procedure and Nei's genetic distances
cladistic biogeography of the Eureptantia 325–336 & Figs 1–2, 338–340
see also ancestral area, BPA, parsimony and three area statements
complementary areas for diversity 129, 139, 147–151 App. 2
development of aquatic insect eggs 223–224, 226–228
see also Bonferroni, Hartley, regression and Tukey tests
egg crypsis in blackbirds 162–163, 167–168
see also CATMOD, Mann-Whitney U- and Wilcoxon signed-rank tests
evenness in moth populations 77–78
first fossil anthomyids 441–442, 445–447
flower-fruit size allometry 401–405
see also statistics
geographic cohesiveness in an intertidal snail 61–64, 68, 70, 72
principal components 61–65 & Tab. 1 & Fig. 2, 67
growth similarities in tits 343–346, 349, 352
see also ANOVAs, bootstraps, correlations, eigenvalues, log-transformations, regressions and variances
monandry in damselflies 86–87, 92–93

- Jolly-Seber Stochastic Method 92
 mtDNA variability and male-lethal symbionts 454,
 461–462 & Figs 2–3, 467
 multivariate genetic differentiation 173–192 & Tab.
 3 & Figs 1 & 3–4
see also ANOVA, Constant Row Total-Multiple
 Correspondence Analysis (CRT-MCA),
 Correspondence (CA), Multiple Correspondence
 (MCA), Multivariate Analysis of Variance
 (MANOVA), Principal Components (PCA) and
 Principal Coordinates (PCoA)
 ornaments are armaments 390
 pyrrolizidine alkaloids in ithomiine butterflies 100,
 103–104, 111–113–114 & Tabs 7–9, 119
see also retention indices
 RAPD variation in *Iliamna* 357, 359–361, 364
see also AMOVA, ANOVA, cluster, HOMOVA
 and statistical analyses
 reliability of synapomorphic analyses 483–497 &
 Figs 1–2 & Tabs 1–3 & Apps 1–2 (& Tabs A1–
 A4)
 Southeast Asian shrews 199, 201–202, 207,
 209–210, 213–215
see also computations, Brook's parsimony,
 Mantel's tests and zoogeography
 specialization and speciation in shrimps
 behavioural 309, 311–312, 317–318
see also binomial and Fisher's exact tests
 genetic 309–310, 312–314 & Fig. 1 & Tab. 1
see also BIOSYS-1, Nei's distances and
 UPGMA
 morphometric 309–311, 314–317 & Tab. 2 &
 Figs 2–3, 320
see also ANOVA, CANDISC, canonical variates
 (CV), Ryan's Q test and Wilks' lambda
see also statistical analyses
 *analysis of molecular variance: RAPD variation in
Iliamna *see* AMOVA
 *ancestral area analysis: cladistic biogeography of the
 Euplantia 325–326, 328–329, 331 & Fig. 1,
 333–336 & Tabs 1–2, 338–339
 ancestry, arthropod 1–2, 41 Fig. 4, 47, 49
 comparison between crustaceans and early
 tracheates 20–22
 groundplan of a polyramous limb 7–10
 problem of radiation 22–24
Andricus quercuscalicis (oak gall wasps): genetics of ice
 ages 260
 anenomes (*Actinia tenebrosa*): intertidal communities 61,
 64 Tab. 1
 angiosperms: sex chromosomes 36
 animals: genetics of ice ages 251–252, 257, 261–263
 anisopteran *Sympetrum danae*: paired spines on accessory
 penes 95
 Annelida: arthropod evolution 25 Fig. 1
Anolis carolinensis (lizards): visual signals 391–392 Tab. 1
Anomalocaris: Cambrian non-arthropod 11, 26
 Anomopoda: arthropod evolution 3
Anopheles maculatus (Culicidae): development of aquatic
 insect eggs 242 App. 2
 Anostraca: arthropod evolution 3, 14, 22, 38
 ANOVAs
 butterfly-ant mutualism 289
 growth similarities in tits 345–346
 multivariate genetic differentiation 179
 RAPD variation in *Iliamna* 359
 specialization and speciation in shrimps 311, 315
 Tab. 2
 Antarctica: development of aquatic insect eggs
 222–223
 Antarcticopleraria (Plecoptera): development of
 aquatic insect eggs 230
 antelopes, Puku (*Kobus vardon*): visual signals 391–392
 Tab. 1
Anthodiscus trifolius: complementary areas for diversity
 148 App. 2
Anthomyia: fossil anthomyids 442
 Anthomyiidae (Diptera): first fossil anthomyid and
 comments on clade definitions 441–451 & Figs
 1–5
 *antipredator behaviour: aggressive nest protection by
 waders 409–413, 426–429
see also predator regimes
 ants: population genetic effects of a butterfly
 mutualism 287–303 & Figs 1–4 & Tabs 1–5
see also *Iridomyrmex*
 anurans: ornaments are armaments 386
see also frogs and toads
Aphriza virgata: aggressive nest defence 438 App.
Apis
A. mellifera: *Eupatorium* flowers 109
 genetics of ice ages 261
A. mellifera (honey bees) 275 App.
 Apocynaceae: pyrrolizidine alkaloids in ithomiine
 butterflies 99, 101, 108–109 Tab. 4 & Fig. 3, 111
see also *Prestonia*
 *aquatic insects: monandry in damselflies 85
Arabidopsis thaliana: inbreeding depression 473
 Arachnida: compound eyes 39, 41 Fig. 4
 Araneidae (*Nephila clavipes*; orb weaving spiders):
 predation of ithomiine butterflies 99–101, 105
 Archosauria: statistics of synapomorphies 489
 arcsine transformations: statistics of synapomorphies
 487, 489, 496 App. 2
 Arctiidae (moths): pyrrolizidine alkaloids 99–100, 114,
 117–118
see also *Hyalurga*, *Scearctia* and *Utetheisa*
 Arctoperlaria (Plecoptera): development of aquatic
 insect eggs 230
see also Systellognatha
 areas: complementary for diversity 125–132, 134–140
 & Fig. 5
 area names 144–157 Apps 1–4
Arenaria: aggressive nest defence
A. interpres 438 App.
A. melanocephala 438 App.
 Argia (Coenagrionidae): development of aquatic insect
 eggs
A. moesta 240 App. 2
A. vivida 240 App. 2
 arginine phosphokinase (EC 2.7.3.3): *Bembicium vittatum*
 (snails)
Apk 1: 62, 65, 67 & Tab. 2, 72
Apk 3: 62, 65, 67 Tab. 2, 72
 *aridity: conservation strategies, stress and diversity
 471, 474–476
 Arizona (Chiricahua Mountains): **Nemoria arizonaria*
 (emerald moths) 279–280
Armadillidium: isopod 35

- armaments: an evolutionary explanation 385–394 & Tab. 1
- Artemia*: branchiopod crustacean 7, 24, 27, 33, 48
- Arthropleura*: arthropod evolution 8
- Arthropoda: reflections on evolution 1–55 & Figs 1–4
- Arthropoda incertae sedis* 31–32 & Fig. 2
- eyelessness in early arthropods 40–44 & Fig. 4
- polyphyletic concepts 25–34 & Figs 1–2
- proto-mandibulates 19–20
- radiation from a common ancestor 22–24
- Asclepiadaceae: pyrrolizidine alkaloids in ithomiine butterflies 106–107 Fig. 2 & Tab. 1, 210 Tab. 5
- Asclepias curassavaria* 106–107 Fig. 2 & Tab. 1, 110 Tab. 5
- Asia
- macaques (molecular data) 276 App.
- Southeast (SE): shrews *Crocodyra* 197–219 & Figs 1–6 & Tabs 1–4 7 App
- assays: pyrrolizidine alkaloids in ithomiine butterflies
- bioassay 99, 105
- colorimetric 104, 111–112 & Tab. 7
- Asteraceae
- flower-fruit size allometry: *Crepis* 401–407 & Figs 1–2
- pyrrolizidine alkaloids 100, 106–110 Figs 2–3 & Tabs 1 & 3–5, 114
- see also
- Adderstoma*, *Bidens*, *Cosmos*, *Ophrystoma*, *Senecio*, *Trichogonia* and *Wulfia*
- Eupatorieae, Heliantheae, Mutisieae and Vernonieae
- Atelocerata: arthropod evolution 2, 4–5
- Atlantic Ocean
- cladistic biogeography of the Eureptantia 340
- Eastern (EAT) 325, 327 Fig. 1, 329, 331–339 & Tabs 1–2 & Figs 1–6
- see also English Channel and Norway:
- Paradrepanophorus*, *Pumettia* and *Uniporus*
- Western (WAT) 325, 327 Fig. 1, 329, 333 Tabs 1–2 & Figs 1–4, 339
- see also Caribbean and North America:
- Curvanemertes* and *Hubrechtomertes*
- genetics of ice ages 249–250, 252, 262
- Atlantic/Gulf coasts 262, 272 App.
- ATP production: energy cost 475
- Atyidae (decapod prawns): arthropod evolution 4
- Australia
- Bembicium vittatum* (snails): Albolhos archipelago 57–59, 64
- collision with Asian volcanic arc 340
- Eureptantia: Great Barrier Reef (Gbr) 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
- genetics of ice ages 249, 257, 260, 276 App.
- monochromatic *Ichnura aurora* females (damselflies) 96
- New South Wales: population genetic effects of a butterfly-ant mutualism 287–288, 290–292 & Fig. 1 & Tab. 1
- Austrocercella* (Notonemouridae): development of aquatic insect eggs
- A. alpina* 239 App. 2
- A. hynesi* 239 App. 2
- A. illiesi* 239 App. 2
- A. tilyardi* 239 App. 2
- Austrocochlea constricta* (trochid): Abrolhos archipelago 61, 63–64 & Tab. 1
- Aves
- statistics of synapomorphies 489–492 & Tabs 2–3, 467–498 App. 2 & Tab. A4
- see also birds
- avocets: aggressive nest defence 422, 427
- see also *Recurvirostra*
- Aysheaia*: Mid-Cambrian Burgess Shales 15
- bacterial gill disease: *Onchorhynchus mykiss* (rainbow trout) 474
- Baetidae (Ephemoptera): development of aquatic insect eggs 240 App. 2
- Baetis rhodani* 240 App. 2
- B. soror* 240 App. 2
- see also *Cloem*
- Balkans: genetics of ice ages 250–252, 259, 262–265 & Fig. 7
- Baltic island of öland: *Crepis tectorum* subsp. *pumila* 402
- Bartramia longicauda* (upland sandpipers): aggressive nest defence 426, 437 App.
- bass, Black Sea (*Centropristis striata*): molecular data 272 App.
- bauplane: arthropod evolution 23, 27
- groundplans 1, 7–10
- beech pollen cores: North America 250–251
- bees (*Apis*)
- A. mellifera*: *Eupatorium* flowers 109
- genetics of ice ages 261
- A. mellifera*: molecular data 275 App.
- beetles
- cerabycid (*Tetraopes tetraophthalmus*): extinction/recolonization 303
- coccinellid: mtDNA variability and male-lethal symbionts 467
- genetics of ice ages 251, 262, 264
- Cicindela* 264
- C. dorsalis* (tiger beetles): molecular data 272 App.
- Belgica antarctica* (chironomid flies): development of aquatic insect eggs 222
- Belize: sponge-dwelling alpheid shrimps 307–309, 312–318 & Figs 1–4 & Tabs 1–2, 320–321
- Bembicium* (snails): geographic cohesiveness 61
- B. vittatum* 57–74 & Figs 1–5 & Tabs 1–4
- Berger-Parker Index: measure of diversity 79–80
- Biblis hyperia* (nymphalines): experimental spider prey 105
- Bidens brasiliensis* (Asteraceae): Ithomiinae visits 110 Tab. 5
- binomial tests: shrimp behavioural responses 312, 317–319 & Fig. 5
- bioassays: pyrrolizidine alkaloids in ithomiine butterflies 99, 105, 115, 117 Tab. 11, 119
- *biodiversity: complementary areas 126–128, 130, 138
- *biodiversity: conservation strategies, stress and diversity 471–473
- *biodiversity: genetics of ice ages 247
- biogeography, cladistic: Eureptantia 325–340 & Figs 1–6 & Tabs 1–3
- BIOSYS-1
- SE Asian shrews 201
- specialization and speciation in shrimps 310
- birch: genetics of ice ages 250
- birds

- genetics of ice ages 251, 262
ornaments are armaments 386–387
 see also black-birds, bobolinks, buntings, cocks, cow-, fowl, greenfinches, grouse, juncos, manakins, martins, partridges, peacocks, pheasants, ptarmigan, snipe, sparrows, tits, warblers and widowbirds
 see also Aves
- Bison bison* (bison): acoustic signals 391–392 Tab. 1
bivalves
 hosts: pea crabs 321
 venerid: effects of competition 478
- blackbirds
 acoustic and visual signals
 red-winged (*Agelaius phoeniceus*) 391–393 & Tab. 1
 yellow-headed (*Xanthocephalus xanthocephalus*) 391–392 Tab. 1
 Icteridae: egg crypsis 159–162 & Fig. 1
 Brewer's (*Euphagus cyanocephalus*) 159–162 & Fig. 1, 164–171 & Tabs 1–3 & Figs 2–4
 red-winged (*Agelaius phoeniceus*) 159–162 & Fig. 1, 164–171 & Tabs 1–3 & Figs 2–4
 yellow-headed (*Xanthocephalus xanthocephalus*) 159–162 & Fig. 1, 164–171 & Tabs 1–3 & Figs 2–4
- blackflies: genetics of ice ages 262, 274 App.
 see also *Simulium*
- Blacksburg, Virginia: RAPD variation in *Iliamna* 358 & Fig. 1
- *blue tits (*Parus coeruleus*): growth similarities 343–354 & Tabs 1–5 & Fig. 1
- bobolinks (*Dolichonyx oryzivorus*): acoustic signals 391–393 & Tab. 1
- *body size (mass): aggressive nest protection by waders 409–412, 414–415, 418–429 & Fig. 3 & Tabs 1–2
- Bombina* (toads): genetics of ice ages 257, 264, 266
 B. bombina (firebellied toads) 263, 275 App.
 B. variegata 263, 275 App.
- Bond cycles: genetics of ice ages 249
- Bonferroni
 development of aquatic insect eggs: adjustment 228
 geographic cohesiveness in an intertidal snail
 correction 67 & Tab. 2
 test 62
- bootstrap procedures
 butterfly-ant mutualism 292, 294, 296–299 & Figs 3–4 & Tab. 5
 growth similarities in tits 346–348 & Tabs 1–2, 351 Tab. 5
- Boraginaceae: pyrrolizidine alkaloids 106–110 Figs 2–3 & Tabs 1 & 3 & 5, 114, 118
 see also *Heliotropium*
- Borneo: SE Asian shrews 197–198, 200 Fig. 1, 202–204 & Fig. 2 & Tab. 1, 208–211 & Figs 5–6, 214, 219 App.
- Borreria latifolia* (Rubiaceae): pyrrolizidine alkaloids in ithomiine butterflies 106–108 & Fig. 2 & Tabs 1 & 3, 110 Tab. 5
- bowfin (*Amia caba*): molecular data 272 App.
- *BPA (Brooks Parsimony Analysis): cladistic biogeography of the Eureptantia 325–342 & Figs 1–6 & Tabs 1–3
- BPA (Brooks Parsimony Analysis): Southeast Asian shrews 197, 199, 202, 208–214 & Figs 3–6
- Branchinecta ferox*: arthropod evolution 3
- Branchiopoda: arthropod evolution 2–4, 13–14, 20, 23, 28–29, 41 Fig. 3
 see also *Artemia*, *Branchinecta* and *Rehbachella*
- Branchiura: arthropod evolution 8, 41–42 & Fig. 4
- Branta canadensis* (Canada geese): molecular data 273 App.
- Brazil
 complementary areas for diversity 125, 134
 San Paulo (Amarais wood): pyrrolizidine alkaloids in ithomiine butterflies 101, 108–110 & Tabs 3 & 5
- Bredocaris*: Cambrian maxillopodan 4, 16, 54–55
- breeding latitudes: aggressive nest protection by waders 409–412, 415, 420–423–425 & Fig. 6 & Tab. 1, 429
- brigalow (*Acacia harpophylla*: host plants): butterfly-ant mutualism 288
- Brinkmannia mediterranea*: eureptantian nemerteans (ME) 327 Fig. 1, 329, 332 Fig. 2
- Britain
 genetics of ice ages 249, 258–259
 see also England
- broken stick (BS) model: evenness in moth populations 75, 77–78 & Fig. 1, 80–82
- Brooks parsimony analyses (BPAs)
 cladistic biogeography of the Eureptantia 325–342 & Figs 1–6 & Tabs 1–3
 Southeast Asian shrews 197, 199, 202, 208–214 & Figs 3–6
- Brugmansia suaveolens* (host plants): *Placidula euryanassa* (ithomiine larvae) 118
- Buchloe dactyloides* (Poaceae): use of AMOVA 360
- budworms, spruce (*Choristoneura*): genetics of ice ages 266, 273 App.
- Bufo*
 acoustic signals: *B. calamita* (natterjack toads) 391–392 Tab. 1
 complex colonization scheme: *B. marinus* 188
- bugs, house martin (*Oeciacus hirundinis*): ectoparasites 474
- buntings, corn: ornaments are armaments 387
- Burgess Shales: arthropod evolution 12, 15, 23, 26, 46
- Burhinus*: aggressive nest defence
 B. oedinemus 436 App.
 B. senegalensis 413, 436 App.
- butterflies
 genetics of ice ages 262, 264, 274 App.
 see also *Evelades*, *Heliconius* and *Neruda*
 ithomiine: pyrrolizidine alkaloids 99–123 & Figs 1–3 & Tabs 1–11
 see also Danainae, Ithomiinae and Nymphalidae
 population genetic effects of ant mutualism 287–305 & Figs 1–5 & Tabs 1–5, 300–301
 see also *Jalmenus*
 sexual size dimorphism 96–97
 swallowtail (Papilionidae): colour polymorphism 278
- Calanoida: arthropod evolution 4, 12
- Caledia*: hybrid zone barriers 259
- Calidris*: aggressive nest defence 422
 C. acuminata 439 App.
 C. alba 438 App.
 C. alpina (dunlin) 417 Fig. 2, 439 App.
 C. bairdi 439 App.
 C. canutus 438 App.

- C. ferruginea* (curlew sandpipers) 426, 439 App.
C. fuscicollis 439 App.
C. himantopus 439 App.
C. maritima 439 App.
C. mauri 439 App.
C. melanotos 439 App.
C. minuta 439 App.
C. minutilla 439 App.
C. ptilocnemis 439 App.
C. pusilla 438 App.
C. ruficollis 439 App.
C. subminuta 439 App.
C. tenuinckii 439 App.
C. tenuirostris 438 App.
- California
 egg crypsis in blackbirds 164, 168
Peromyscus maniculatus (deer mouse) 274 App.
 callimorphine: pyrrolizidine alkaloid 117
Calopogonium sp. (Leguminosae): *Aeria olena* visits
 106–107 Fig. 2 & Tab. 1
Calythea: systematic relationships 445–446
 Cambrian: arthropod evolution 3–4, 9–16, 18–20,
 22–32 & Figs 1–2, 42–47
 convergence in compound eyes 37–38, 40
 ‘explosion’ 46, 54–55
 eyeless Skaracarida 41
 Pre-Cambrian 10, 13–14, 20, 25–27, 33
 (late): Vendian 26
 problematica 41 Fig. 4
see also Anomalocaris, Cambropodus, Canadaspis,
Rehbachella, Sanctacaris and Sirotoecerus
Cambropodus: arthropod evolution 9, 13, 19–20, 32
 *camouflage: egg crypsis in blackbirds 159
 Canada
Crepis tectum 403
 damselflies: pruinescence 94
 genetics of ice ages 250–251, 272–273 App.
Canadaspis: Cambrian arthropod 8, 26
 CANDISC procedure of SAS: specialization and
 speciation in shrimps 311
Canis
 molecular data
C. latrans (coyotes) 273 App.
C. lupus (wolves) 273 App.
 placental dogs: convergence 35
 canonical variate (CV) analysis: specialization and
 speciation in shrimps 310–311, 314, 316–317
 Figs 2–3
 *capillary gas chromatography: pyrrolizidine alkaloids
 (PAs) in ithomine butterflies 99, 104–105
 Capniidae (Euholognatha): development of aquatic
 insect eggs 238 App. 2
Capnia atria 238 App. 2
C. bifrons 238 App. 2
C. gracilaria 238 App. 2
C. nana 238 App. 2
see also Mesocapnia and Utacapnia
 Carboniferous: arthropod evolution 8–9, 30, 33, 43
see also Arthropleura
 Caribbean
 sponge-dwelling alpheid shrimps 307–309
see also Belize and Panama
 Western Atlantic Ocean (WAT): Eureptantia 327
 Fig. 1, 329, 335 Fig. 4
 caribou: ornaments are armaments 387
Cariniana: complementary areas for diversity
C. asterophora 150 App. 2
C. asterotricha 150 App. 2
C. crenatum 150 App. 2
C. ianirensis 147 App. 2
C. kuhlmannii 150 App. 2
C. pachyantha 149 App. 2
C. pauciramosa 148 App. 2
C. pendulifera 148 App. 2
C. pyramidata 147 App. 2
C. rubra 148 App. 2
C. uapensis 150 App. 2
 Carpathians: genetics of ice ages 264–265 & Fig. 7
 Caryocaraceae: complementary areas for diversity
 125, 128–129, 132–138 & Figs 1 & 3–4 & 6–7
Caryocar barbinervi 150 App. 2
C. coriaceum 148 App. 2
C. costaricense 147 App. 2
C. harlingei 148 App. 2
 Caspian Sea: genetics of ice ages 252–253
Casterosteus aculeatus (three-spined sticklebacks):
 ornaments are armaments 388, 391–392 Tab. 1
 *caterpillars: *Nemoria arizonaria* (emerald moths)
 277–283 & Tab. 1 & Fig. 2
 catkins, oak: effect of light and diet on a polymorphic
 caterpillar 277–283 & Tab. 1 & Fig. 2
 CATMOD procedure: egg crypsis in blackbirds 167
Catoptrophorus semipalmatus: aggressive nest defence 438
 App.
 Catstat: geographic cohesiveness in an intertidal snail
 62
 Caucasus: genetics of ice ages 249, 251–253, 262
Centropomus striata (Black Sea bass): molecular data 272
 App.
 Cephalocarida: arthropod evolution 4, 40–41 & Fig.
 4, 55
 cerabycid beetles (*Tetraopes tetraopthalms*): extinction/
 recolonization 303
Cervus elaphus (red deer): acoustic and visual signals
 391–392 Tab. 1
Cestrum sp. (Solanaceae): *Tithorea harmonia* visits
 108–109 Tab. 4 & Fig. 3
 chaffinches: open-nesting 344
 character analysis: first fossil anthomyid 441, 445–447
 Charadriiformes: aggressive nest protection by waders
 410, 418
 Charadriomorpha 427
see also avocets, crab plover, gulls, lapwings,
 oystercatchers, plovers, pratincoles, seedsnipe,
 sheathbills and thick-knees
Charadrius: aggressive nest defence
C. alexandrinus 437 App.
C. asiaticus 437 App.
C. dubius 437 App.
C. hiaticula 436 App.
C. leschenaultii 437 App.
C. melodus 437 App.
C. mongolus 437 App.
C. montanus 437 App.
C. morinellus (dotterels) 428, 437 App.
C. pecuarius 437 App.
C. placidus 437 App.
C. semipalmatus 437 App.
C. veredus 437 App.
C. vociferus 437 App.

- C. wilsonia* 437 App.
cheatgrass: spread in North America 251
cheetahs (*Acinonyx jubatus*): genetic stress 472–473
Chelicerata: arthropod evolution 5–6, 8, 23–25 & Fig. 1, 41–42 & Fig. 4
Chelonia: statistics of synapomorphies 490–492 & Tabs 2–3
*chemical defence: pyrrolizidine alkaloids (PAs) in ithomiine butterflies 99, 118–120
Chen p. oerulescens (snow geese): molecular data 273 App.
Chengjiang Deposits: arthropod evolution 12, 15, 46
chi-square tests
aggressive nest protection by waders 418–423
geographic cohesiveness in an intertidal snail 62, 65–66 & Fig. 3
chickadees, North American (*Parus*): genetics of ice ages 261, 264
P. atricapillus: molecular data 273 App.
P. carolinensis 273 App.
Chilopoda: arthropod evolution 6, 41 Fig. 4, 43
Chironomidae (Diptera): development of aquatic insect eggs 231–232
Chiroxiphia linearis (swallow-tailed manakins): visual signals 391–392 Tab. 1
Chloroperlidae (Systellognatha): development of aquatic insect eggs 240 pp. 2
Chloroperla tripunctata 230, 240 App. 2
Choristoneura (spruce budworms): genetics of ice ages 266, 273 App.
Chorthippus (grasshoppers): genetics of ice ages 247, 261, 266
C. parallelus 247, 251, 258–265 & Figs 5–7, 275 App.
C. p. erythropus: Iberia 258, 263, 266
C. p. parallelus: Britain to Siberia and Finland to Turkey 258, 263, 266
C. p. tenuis: Greece 258
chromatography: pyrrolizidine alkaloids (PAs) in ithomiine butterflies 99, 104–105
Chrysiptera cyanea (damsel fish): visual signals 391–392 Tab. 1
Chrysobalanaceae: complementary areas for diversity 125, 128–129, 132–138 & Figs 1–4 & 6–7
Chrysobalanus cuspidatus 148 App. 2
C. venezuelanus 150 App. 2
cicadas, periodic (*Magiciada*): molecular data 274 App.
cichlid fish: genetics of ice ages 262, 264
mbuna: molecular data
Melanochromis auratus 275 App.
M. heterochromis 275 App.
Pseudotropheus zebra 275 App.
Cicindela (beetles): genetics of ice ages 264
C. dorsalis (tiger beetles): molecular data 272 App.
Cirrropedia: eyeless adults 42
Citellus beecheyi (California ground squirrels): blackbird egg predation 162
citrine forktails (*Ischnura hastatum*): monandry in damselflies 87
CK-1 and CK-2 (EC 2.7.3.2): Southeast Asian shrews 200
CK-1 204 Tab. 1, 206 Tab. 3
CK-1 and *CK-2* 205 Tab. 2
CK-2 204 Tab. 1, 206 Tab. 3
*cladistic biogeography: Euplantia 325–340 & Figs 1–6 & Tabs 1–3
*cladistics: synapomorphies 483–497 & Figs 1–2 & Tabs 1–3 & Apps 1–2 (& Tabs A1–A4)
cladistics
arthropod evolution 5–6, 12, 23–24, 39, 48–49, 55
first fossil anthomyid and comments on definitions 441, 445–451 & Figs 3–5
*cladograms: synapomorphies 483–486 & Figs 1–2
clams, coot (*Mulinia lateralis*): heterozygosity 174
classification: first fossil anthomyid and comments on clade definitions 441–442, 447–451 & Figs 4–5
Clethrionomys: hybrid zone barriers 259
climates: genetics of ice ages 248–253, 255–256, 259–260 & Fig. 6, 262–263
see also palaeoclimates
Cloeon triangulifer (Bactidae): development of aquatic insect eggs 240 App. 2
cluster analysis: RAPD variation in *Ilamna* 357, 365
Coccinia: large unicorneal eyes 39
cockroaches (*Naupeta cinerea*): ornaments are armaments 388
cocks of the rock: ornaments are armaments 389
Coella tiuensis: eurypteric nemertean (EIO: Tiur) 327–328 & Fig. 1, 332 Fig. 2, 339
Coenagrionidae (Odonata): development of aquatic insect eggs 223, 240 App. 2
Coenagrion puella 240 App. 2
see also *Argia*, *Enallagma* and *Ischnura*
Coenosopsia: first fossil anthomyid and comments on clade definitions 441, 445–448 & Fig. 3, 450–451 & Fig. 5
Coenosopsites (gen. nov.): first fossil anthomyid 442, 446, 450
C. poinari (sp. nov.) 441–451 & Figs 1–5
Coffea (Rubiaceae): ithomiine visits 110 Tab. 5
Colaptes auratus (woodpeckers): genetics of ice ages 263
Coleomegilla maculata: mtDNA variability and male-lethal symbionts 467
collecting and scoring: evenness in moth populations 76
Collembola: eyeless or without compound eyes 41 Fig. 4, 43
Coloburiscidae (Ephemeroptera): development of aquatic insect eggs 240 App. 2
Coloburiscoides nr haleuticus 240 App. 2
coloniality: aggressive nest protection by waders 409, 411, 417, 422–426 & Tabs 1–2, 428, 436–439 App.
colonization: genetics of ice ages 247, 252–254 & Fig. 1, 259–261 & Fig. 6, 264, 266
coloration
crisis in blackbirds' eggs 159–171 & Figs 1–4 & Tabs 1–3
monandry in damselflies 85, 91–94 & Tab. 1, 96
colorimetric assay: pyrrolizidine alkaloids in ithomiine butterflies 104, 111–112 & Tab. 7
*colour change: *Nemoria arizonaria* (emerald moths) 277–283 & Figs 1–2 & Tab. 1
Columbia livia (feral rock doves): parasitic stress 474
*competition: conservation strategies, stress and diversity 471, 473, 477–479
*competition: evenness in moth populations 75, 77, 82
*complementarity: areas for diversity 126–128, 131–132, 135
Component 2.0: (computer program): cladistic biogeography of the Euplantia 328, 336, 340

- *compound eyes: arthropod evolution 1–2, 19, 37–40, 54
- computations
- arthropod evolution: cladistics 23–24
 - cladistic biogeography of the Eurentantia 336
 - see also* bootstrap values, Component 2.0, DELTRAN, MacClade 3.0 and PAUP
 - complementary areas for diversity 126, 130–131, 139
 - see also* WORLDMAP software
 - geographic cohesiveness in an intertidal snail Catstat 62
 - Statview SE + Graphics 61
 - Systat version 5.2: 63
 - mtDNA variability and male-lethal symbionts 462
 - multivariate genetic differentiation
 - ADE 192
 - FSTAT 181
 - SE Asian shrews 197, 201
 - see also* BIOSYS-1, PAUP and 'R' programs
 - simulations
 - genetics of ice ages 252–255 & Figs 1–2
 - statistics of synapomorphies 487, 492–493, 496–498 App. 2 & Tab. A4
 - specialization and speciation in shrimps *see also* BIOSYS-1 and CANDISC
- conclusions
- conservation strategies, stress and diversity 479–480
 - development of aquatic insect eggs 233–234
 - mtDNA variability and male-lethal symbionts 468
 - multivariate genetic differentiation 192
- *conservation: complementary areas for diversity 125–130, 132, 138–139
- Conservation International (CI) 127, 129
- conservation: strategies, stress and diversity 471–473, 476–479
- *conservation genetics: RAPD variation in *Iliamna* 357
- constant row total-multiple correspondence analysis
- (CRT-MCA): multivariate genetic differentiation 173, 176–184 & Fig. 1 & Tabs 1–3, 186–192 & Fig. 6 & Tabs 4–5
- constraints: flower-fruit size allometry 401–402, 406
- contact guarding: monandry in damselflies 85
- contingency tests: geographic cohesiveness in an intertidal snail 62, 66–67 Fig. 3 & Tab. 2
- *convergence: arthropod evolution 1, 18, 34–37, 47
- compound eyes 37–40
- Copepoda: arthropod evolution 4–5, 8, 12, 15, 41 Fig. 4
- copper: experimental stress on freshwater plankton 479
- copulations: monandry in damselflies 85–91 & Tab. 1, 95–96
- *coral reefs: specialization and speciation in shrimps 307–309, 320–321
- coral-dwelling mussels 321
 - see also* Agelas, Hyattella, Hymeniacidon, Lissodendoryx, Oceanapia and Xestospongia
- Corduliidae (Odonata): development of aquatic insect eggs 240 App. 2
- Cordulia shurtleffi* 240 App. 2
- Coregonus: genetics of ice ages 262
- C. dupeaformis* (lake whitefish) 261, 272 App.
- correlations
- aggressive nest protection 424, 426, 428
 - Spearman's Rank 421–423
 - development of aquatic insect eggs 222, 225, 232
 - egg crypsis in blackbirds 159, 166 & Fig. 3
 - flower-fruit size allometry 401–405
 - geographic cohesiveness in an intertidal snail 64
 - Tab. 1, 67–69
 - growth similarities in tits 346–347 & Tab. 1, 350, 353 354
 - multivariate genetic differentiation 176, 178–179 & Tab. 1, 183–185 & Tab. 3 & Fig. 4, 187–188
 - Southeast Asian shrews 197, 202, 206–208, 210 Fig. 3, 215–216
- Correspondence Analysis (CA): population genetics 174–176, 179, 189, 191 & Tab. 4
- corvids
- Corvus* (crows): hybrid zones 257
 - nest predators of waders 416–417, 422, 436–439 App.
- Corythophora amapaensis*: complementary areas for diversity 148 App. 2
- Cosmos bipinnatus* (Asteraceae): Ithomiinae visits 110 Tab. 5
- Costa e Silva wood (San Paulo, Brazil): pyrrolizidine alkaloids in ithomiine butterflies 101, 105–107 & Fig. 2 & Tabs 1–2, 109–110 & Tab. 5, 112, 114
- Couepia*: complementary areas for diversity
- C. amarulae* 148 App. 2
 - C. caraulae* 148 App. 2
 - C. caryoph. s. glabra* 150 App. 2
 - C. cidana* 151 App. 2
 - C. coarctata* 149 App. 2
 - C. exflexa* 147 App. 2
 - C. froesei* 150 App. 2
 - C. glabra* 148 App. 2
 - C. impressa s. cabraliae* 147 App. 2
 - C. insignis* 147 App. 2
 - C. kruckii* 148 App. 2
 - C. latifolia* 149 App. 2
 - C. longipetiolata* 147 App. 2
 - C. maguirei* 148 App. 2
 - C. marleneae* 149 App. 2
 - C. martinii* 147 App. 2
 - C. meridionalis* 151 App. 2
 - C. monteclarensis* 149 App. 2
 - C. morii* 148 App. 2
 - C. multiflora* 151 App. 2
 - C. nultans* 148 App. 2
 - C. paraensis s. cerrad.* 151 App. 2
 - C. parvifolia* 150 App. 2
 - C. pernambucensis* 149 App. 2
 - C. recurva* 149 App. 2
 - C. reflexa* 150 App. 2
 - C. scotmorii* 147 App. 2
 - C. steyermarkii* 147 App. 2
 - C. stipularis* 148 App. 2
 - C. trapeziana* 149 App. 2
 - C. uiti* 148 App. 2
- Cowatari*: complementary areas for diversity
- C. asterophora* 150 App. 2
 - C. asterotricha* 150 App. 2
 - C. calycina* 148 App. 2
 - C. prancei* 149 App. 2
 - C. pyramidata* 147 App. 2
 - C. sandwithii* 149 App. 2
- covariances: statistics of synapomorphies 484, 491, 493

- covariation: monandry in damselflies 85, 91 Tab. 1, 94–95
- cowbirds, brown-headed (*Molothrus ater*): ornaments are armaments 390, 391–392 Tab. 1
- coxa: arthropod evolution 1–5, 15, 32
- crabs
- fiddler (*Uca pugnator*): ornaments are armaments 386, 391–392 Tab. 1
 - hermit: hydroids encrusting shells 321
 - horseshoe (*Limulus polyphemus*): molecular data 272 App.
 - pea: bivalve hosts 321
- Crassostrea virginica* (American oysters): molecular data 272 App.
- Crepis* (Asteraceae): flower-fruit size allometry 401–407 & Figs 1–2
- C. tectorum* 401–406 & Fig. 2
 - subsp. *pumila* 401–404 & Fig. 4, 406
- Cretaceous: eurentatic phylogenetic trees 339
- crickets
- cave (*Dolichopoda*): genetics of ice ages 261
 - D. laetitiae*: molecular data 275 App.
 - gregarious (*Ampicacusta maya*): acoustic signals 391–392 Tab. 1
- *Crocodyrines: SE Asian shrews 197
- Crocodyr* 197, 199, 203–206 & Tabs 1–3, 213–216
- C. attenuata* 215
 - C. batuensis* 214
 - C. beatus* 203–204 & Tab. 1, 208–215 & Tab. 4 & Figs 3–5, 219 App.
 - C. beccarii* 203, 205 Tab. 2, 207–212 & Tab. 4 & Figs 3–5, 215, 219 App.
 - C. brunnea* 205 Tab. 2, 208–212 Tab. 4 & Figs 3–5
 - C. b. brunnea* 219 App.
 - C. brunnea-C. orientalis* clade 211
 - C. elongata* 203, 206 Tab. 3, 208–213 & Tab. 4 & Figs 3–5, 219 App.
 - C. foetida* 203, 207, 210, 213–214
 - C. foetida* ssp. 212 Fig. 5
 - f. doriae* 203–204 & Tab. 1, 206–211 & Tab. 4 & Figs 3–4, 219 App.
 - f. foetida* 204 Tab. 1, 208–211 Tab. 4 & Figs 3–4, 219 App.
 - f. kelabit* 204 Tab. 1, 208–211 Tab. 4 & Figs 3–4, 219 App.
 - C. fuliginosa* 204 Tab. 1, 207–215 & Tab. 4 & Figs 3–6, 219 App.
 - C. grandis* 214
 - C. grayi* 203–204 & Tab. 1, 208–210 & Tab. 4 & Fig. 3, 212–215 & Fig. 5, 219 App.
 - C. grayi-C. beatus* cluster 207
 - C. hutani* 204 Tab. 1, 207–212 & Tab. 4 & Figs 3–5, 215, 219 App.
 - C. lea* 206–212 & Tabs 3–4 & Figs 3–5, 219 App.
 - C. lepidura* 205 Tab. 2, 207–212 & Tab. 4 & Figs 3–5, 214–215, 219 App.
 - C. levicula* 206 Tab. 3, 208–213 & Tab. 4 & Figs 3–5, 219 App.
 - C. malayana* 203, 211–212 & Fig. 5, 219 App.
 - malayana 1* 204 Tab. 1, 208–211 Tab. 4 & Figs 3–4
 - malayana 3* 204 Tab. 1, 208–211 Tab. 4 & Figs 3–4
 - C. malayana-C. negligens* cluster 207
 - C. mindorus* 203–204 & Tab. 1, 206, 208–214 & Tab. 4 & Figs 3–5, 219 App.
 - C. monticola* 205 Tab. 2, 207–215 & Tab. 4 & Figs 3–6, 219 App.
 - C. musseri* 206 Tab. 3, 208–213 & Tab. 4 & Figs 3–5, 219 App.
 - C. musseri-C. elongata-C. levicula* cluster 207
 - C. negligens* 204 Tab. 1, 208–212 & Tab. 4 & Figs 3–5, 219 App.
 - C. negrina* 214
 - C. nigripes* 197, 203, 207, 210, 212 Fig. 6, 214
 - C. nigripes* ssp. 212 Fig. 5
 - C. n. lipara* 206 Tab. 3, 208–211 Tab. 4 & Figs 3–4, 213, 219 App.
 - C. n. nigripes* 203, 206 Tab. 3, 208–211 Tab. 4 & Figs 3–4, 219 App.
 - C. nigripes-C. foetida* cluster 207
 - C. nigripes-C. malayana* clade 215
 - C. olivieri* 199, 201, 203–204 Tab. 1, 208–211 Tab. 4 & Figs 3–4
 - o. lauwana* 203, 205 Tab. 2, 208–211 Tab. 4 & Figs 3–4, 219 App.
 - o. orientalis* 203, 205 Tab. 2, 208–211 Tab. 4 & Figs 3–4, 219 App.
 - C. orientalis*
 - C. orientalis-C. brunnea* clade 207
 - C. ssp.* 212 Fig. 5
 - C. palawanensis* 214
 - C. paradoxura* 205 Tab. 2, 207–212 & Tab. 4 & Figs 3–5, 215, 219 App.
 - C. rhoditis* 206–213 & Tabs 3–4 & Figs 3–5, 219 App.
- Crocodylia: statistics of synapomorphies 489–492 & Tabs 2–3, 497–498, App. 2 & Tab. A4
- Crotalaria* (Fabaceae): pyrrolizidine alkaloids 100, 118
- Croton lundianus* (Euphorbiaceae): ithomiine visits 108 Tab. 3, 110 Tab. 5
- crows (*Corvus*): hybrid zones 257
- Crustacea
- arthropod evolution 1–31 & Figs 1–2
 - ancestors compared with early tracheates 20–22
 - compound eyes 37–40
 - epilogue 49
 - eyeless ancestors 40–41 & Fig. 4
 - monophyly 33–36 & Fig. 3
 - Ur-Crustacea 20, 40–41 & Fig. 4
 - sponge-dwelling alpheid shrimps 307–308, 320
 - crypsis: eggs in blackbirds 159–160, 165, 169–171
 - quantification 162–163
 - see also patterns and survival
- *crypsis: *Nemoria arizonaria* (emerald moths) 277
- Culicidae (Diptera): development of aquatic insect
- eggs 221, 225, 228, 230–232 & Fig. 5, 242 App. 2
 - Culex anulisrostris* 242 App. 2
 - C. pipiens* 242 App. 2
 - C. pipiens fasciatus* 232
 - C. restuans* 242 App. 2
 - see also *Aedes*, *Anopheles* and *Toxorhynchites*
- Cupienius gelazi* (wandering spiders): acoustic signals 391–392 Tab. 1
- curlews: aggressive nest defence 427
- see also *Numenius*
- Curranemertes natans*: eurentatic nemertean (WAT: Caribbean) 327 Fig. 1, 329, 332 Fig. 2, 339

- Cursorius cursor*: aggressive nest defence 436 App.
- Cyclopoida: arthropod evolution 4
- cyotes (*Canis latrans*): molecular data 273 App.
- Cyprinodon pecosensis* (Pecos pupfish): visual signals 391–392 Tab. 1
- Cyrtodopsis whitei* (stalk-eyed flies): visual signals 391–392 Tab. 1
- *cytoplasmic symbionts: mtDNA variability and male-lethal symbionts 453–454, 457–459, 468
see also *Wolbachia*
- Dactylostemon estrellensis* (Euphorbiaceae): Ithomiinae visits 108 Tab. 3, 110 Tab. 5
- damselfish (*Chrysiptera cyanea*): visual signals 391–392 Tab. 1
- damselflies: evolution of monandry 85–98 & Fig. 1 & Tab. 1
- Danainae (ithomiine butterflies): pyrrolizidine alkaloids 99–100, 109, 119
- Darwinism Evolving. Systematics Dynamics and the Genealogy of Natural Selection*, by D. J. Depew and B. H. Weber. (Book Review: M. F. Claridge) 499–500
- complementary areas for diversity 126, 128–129, 132–134 & Figs 1–4 & 1ts 1–2, 136–139
- nested: statistics of synapomorphies 484–486 & Figs 1–2
- Decapoda
- divergent radiation 28–29
- sponge-dwelling alpheid shrimps 307–308, 314
- deer
- red (*Cervus elaphus*): acoustic and visual signals 391–392 Tab. 1
- white-tailed (*Odocoileus virginianus*): molecular data 274 App.
- deermice (*Peromyscus maniculatus*): molecular data 274 App.
- definitions
- aggressive nest protection by waders 409, 413–414
- first fossil anthomyid and comments on clade definitions 441, 447–451 & Figs 4–5
- Delichon urbica* (house martin): parasitic stress 474
- DELTRAN (program): Eupreptantia 334 Fig. 3, 338
- demographics: monandry in damselflies 90–93 & Tab. 1
- Depew, D. J. and Weber, B. H. *Darwinism Evolving. Systematics Dynamics and the Genealogy of Natural Selection*. (Book Review: M. F. Claridge) 499–500
- detectability on nest: protection by waders 409, 411–412, 415–416, 422, 424–426 & Tabs 1–2, 428, 436–439 App.
- development: aquatic insect eggs 221–233 & Figs 1–5, 236–242 Apps 1–2
- Devon coast (England): *Nucella lapillus* (dogwhelks): multivariate genetic differentiation 173, 175, 180–182 & Fig. 2, 187
- Devonian: arthropod evolution 3–4, 13, 30, 32, 43
- *diapause: development of aquatic insect eggs 221, 223, 225–233 & Fig. 2
- Diaptomus castor*: arthropod evolution 4
- Dicharpetalaceae: complementary areas for diversity 125, 128–129, 132–138 & Figs 1 & 3–4 & 6–7
- Dicharpetalum*: complementary areas for diversity
- D. asplundeanum* 150 App. 2
- D. bernalii* 149 App. 2
- D. costaricense* 150 App. 2
- D. gentryi* 149 App. 2
- D. hammelii* 147 App. 2
- D. pauper* 149 App. 2
- D. steyermarkii* 151 App. 2
- D. stipulatum* 150 App. 2
- Dicliptera pohlana* (Acanthaceae): *Tithorea harmonia* visits 108–109 Tab. 4 & Fig. 3
- *diet-induced polymorphism: *Nemoria arizonaria* (emerald moths) 277–281 & Tab. 1, 283
- Dinocraus cephalotes* (Perlidae): development of aquatic insect eggs 239 App. 2
- Dinopis*: large simple eyes 39
- Diplopoda: arthropod evolution 6–8, 41 Fig. 4
- Diplura: eyeless or without compound eyes 41 Fig. 4, 43, 49
- *Diptera: development of aquatic insect eggs 221, 226–229 & Figs 3–4, 232–233, 242 App. 2
see also Chironomidae, Culicidae, Sciomyzidae and Tipulidae
- Diptera
- Anthomyiidae: first fossil anthomyid and comments on clade definitions 441–451 & Figs 1–5
see also Fanniidae and Muscidae
- Heliotropium transalpinum* flowers 109
- origins of prolegs 34
- Dircenna dero*: Ithomiinae plant visitors 110 Tab. 5
- discussions
- aggressive nest protection by waders 425–429
- butterfly-ant mutualism 297–303 & Figs 3–5
- cladistic biogeography of the Eupreptantia 334–341 & Figs 3–6 & Tab. 3
- complementary areas for diversity 136–140
- development of aquatic insect eggs 229–232 & Figs 4–5
- effect of light and diet on a polymorphic caterpillar 282–283
- egg crypsis in blackbirds 168–171
- evenness in moth populations 81–83
- first fossil anthomyid and comments on clade definitions 447–451
- flower-fruit size allometry 405–406
- geographic cohesiveness in an intertidal snail 70–73
- growth similarities in tits 351–354 & Tabs 5–6
- monandry in damselflies 94–97
- mtDNA variability and male-lethal symbionts 164–168 & Figs 5–6 & Tabs 1–3
- multivariate genetic differentiation 187–191 & Fig. 7 & Tab. 4
- ornaments are armaments 392–394
- pyrrolizidine alkaloids in ithomiine butterflies 115–121 & Tabs 10–11
- RAPD variation in *Itania* 365–367
- SE Asian shrews 213–217
- specialization and speciation in shrimps 319–322
- statistics of synapomorphies 484, 491–493
- *dispersal: genetics of ice ages 247, 251, 253–257 & Figs 2–4
- *dispersal: population genetic effects of a butterfly-ant mutualism 287–289
- Distal-less (Dll)*: limb-patterning gene 7, 29
- distance-Wagner procedure: SE Asian shrews 201, 206–207
- Diura bicaudata* (Perlodidae): development of aquatic insect eggs 228, 230, 232, 239 App. 2

- divergence: genetics of ice ages 247, 267–258, 261–266 & Fig. 7, 272–276 App.
- *diversity: evenness in moth populations 75–77, 79–83 & Fig. 3 & Tab. 1
see also Berger-Parker, Simpson's Indices and log-series
- *diversity: specialization and speciation in shrimps 307–308, 320
- diversity
 complementary areas 126, 128–132, 134, 136–139
 genetic: conservation strategies and stress 471–482
- DNA
 changes: arthropod evolution 47
 DNA-DNA hybridization: Cambrian organisms 29
 RAPD variation in *Iliamna* 357, 359–361
- *DNA sequences: genetics of ice ages 247, 257–262 & Fig. 5, 266
 mtDNA 258–259, 261, 264, 266
- dogs (*Canis*): convergence 35
- dogwhelks (*Nucella lapillus*): multivariate genetic differentiation 173, 175, 180–182 & Fig. 2, 187–190
- dolichoderine ants: population genetic effects of a butterfly mutualism 290
see also *Iridomyrmex*
- Dolichonyx oryzivorus* (bobolinks): acoustic signals 391–393 & Tab. 1
- Dolichopoda* (cave crickets): genetics of ice ages 261
- D. laetitiae*: molecular data 275 App.
- *Dominican amber: first fossil anthomyid 441–443 & Fig. 1
- dotterels (*Charadrius morinellus*): aggressive nest defence 428, 437 App.
- doves
 collared: rapid expansion 251
 feral rock (*Columba livia*): parasitic stress (lice) 474
- dowitchers: aggressive nest defence 427
see also *Limnodromus*
- dragonflies
 development of aquatic insect eggs *see* Odonata
- Libellula luctuosa*: visual signals 391–392 Tab. 1
- Drepanobanda trilineata*: eureptantian nemerteans (EIO: Timor) 327–328 & Fig. 1, 332 Fig. 2
- Drepanogigas albolineatus*: eureptantian nemerteans (ME) 327 Fig. 1, 329, 332 Fig. 2
- Drepanophorella sebae*: eureptantian nemerteans (EIO: Savu) 327–328 & Fig. 1, 332 Fig. 2, 339
- Drepanophorella*: eureptantian nemerteans
D. lifuensis (WP: New Caledonia) 327–329 & Fig. 1, 332 Fig. 2
D. rosea (WIO: Maldives) 327–328 & Fig. 1, 332 Fig. 2, 339
- Drepanophoria pajungae*: eureptantian nemerteans (EIO: Sulawesi) 327–328 & Fig. 1, 332 Fig. 2, 339
- Drepanophorina*: eureptantian nemerteans
D. argus (EIO: Timor) 327–328 & Fig. 1, 332 Fig. 2
D. guineensis (WP: New Guinea) 327–329 & Fig. 1, 332 Fig. 2
D. lata (EIO: Ambon) 327–328 & Fig. 1, 332 Fig. 2
D. savuensis (EIO: Savu) 327–328 & Fig. 1, 332 Fig. 2
- Drepanophoringia paupensis*: eureptantian nemerteans (EIO: Sumba) 327–328 & Fig. 1, 333 Fig. 2, 339
- Drepanophorus*: eureptantian nemerteans 339
- D. modestus* (EIO: Flores) 327–328 & Fig. 1, 332 Fig. 2
- D. rubrostriatus* (ME) 327 Fig. 1, 329, 332 Fig. 2, 339
- Dromas ardeola*: aggressive nest defence 436 App.
- Drosophila* (fruitflies)
 arthropod evolution 21–22, 27–29, 37, 46
 role of the *Distal-less* gene 7
 conservation strategies, stress and diversity
D. buzzati 473, 478
D. melanogaster 473, 477
 genetics of ice ages 266
D. melanogaster: response to desiccation 263
D. montium: molecular data 276 App.
 inversion polymorphism: Principal Components Analysis (PCA) 174
 mtDNA variability 459, 467
D. posttritana 467
D. simulans 454
D. willistoni 467
 ornaments are armaments 388
- Dryas iulia* (Nymphalinae: Heliconiini): non-pyrrolizidine alkaloid insects 109
- Dugesia*: arthropod evolution 45
- dunlin (*Calidris alpina*): aggressive nest defence 417 Fig. 2, 439 App.
- earwigs, common (*Forficula auricularia*): visual signals 391–392 Tab. 1
- Easter Group (Houtman Albrolhous Islands, Western Australia): *Bembicium vittatum* (snails) 60 Fig. 1, 63, 65 & Fig. 2, 67–71 & Tab. 2 & 4 & Fig. 4
- Ecdyonuridae (Ephemoptera): development of aquatic insect eggs 241 App. 2
- Ecdyonurus dispar* 241 App. 2
- E. insignis* 241 App. 2
- E. picteti* 241 App. 2
- E. torrentis* 241 App. 2
- E. venosus* 241 App. 2
- echinatine (pyrrolizidine alkaloid): ithomiine butterflies 99, 102 Fig. 1, 104, 109–110, 113–114, 117, 119
- Echinocloa microstachya* (North American barnyard grass): genetics of ice ages 260
- Echinometra mathaei* (urchins): intertidal communities 61, 64 Tab. 1
- Echitoideae
 Parsonsiaeae: isolation of parsonsine 110
- Prestonia acutifolia*: foodplant of *Tithorea harmonia* larvae 99, 101
- *ecological specialization: sponge-dwelling alpheid shrimps 307, 319–321
- Ecological versatility and Community Ecology*, by Ralph C. Mac Nally (Book Review: Mark O. Hill) 245
- Eemian interglacial: genetics of ice ages 249
- *egg coloration: crypsis in blackbirds 159–171 & Figs 1–4 & Tabs 1–3
see also domestic fowl and village weavers
- *egg predation: crypsis in blackbirds 159–162 & Fig. 1
see also snakes and squirrels
- eggs, aquatic insect: development 221–233 & Figs 1–5, 236–242 Apps 1–2
- eigenvalues
 Constant Row Total-Multiple Correspondence Analysis (CRT-MCA) 178–179, 181–182 & Tab. 2, 186–188 & Fig. 6, 194–195 App.
 growth similarities in tits 346–351 & Tabs 1–2 & 5

- Eisenia foetida* (manure worms): heterozygote advantage 474
- electrophoreses
- development of aquatic insect eggs 232–233
 - geographic cohesiveness in an intertidal snail 62
 - loci and stress 475–476
 - multivariate genetic differentiation 185 Fig. 4
 - population genetic effects of a butterfly-ant mutualism 287, 290–291
 - SE Asian shrews 197, 199–201, 219 App.
 - sponge-dwelling alpheid shrimps 307–310
- Elephantopus mollis* (Vernoniaeae; Asteraceae): *Aeria olena* visits 106–108 Fig. 2 & Tabs 1 & 3, 110 Tab. 5
- elephants (Proboscidae): convergence 36
- Emilia sonchifolia* (Heliantheae; Asteraceae):
- pyrrolizidine alkaloids 106–107 Fig. 2 & Tab. 1, 113–114 & Tab. 9, 117
- emiline: pyrrolizidine alkaloid 114
- Enallagma* (Coenagrionidae): development of aquatic insect eggs
- E. boreale* 240 App. 2
 - E. cyathigerum* 240 App. 2
 - E. ebrium* 240 App. 2
 - E. hageni* 240 App. 2
 - E. yemale* 240 App. 2
- Enchenopa binotata* (treehoppers): genetics of ice ages 266
- *endangered species: RAPD variation in *Iliamna* 357, 360
- endemism: complementary areas for diversity 126–127, 130, 132, 136–137
- see also narrow endemism
- *energy cost: conservation strategies, stress and diversity 471, 473–479
- England
- Devon coast: *Nucella lapillus* (dogwhelks): multivariate genetic differentiation 173, 175, 180–182 & Fig. 2, 187
 - genetics of ice ages 251
 - Woodchester Park Field Centre: evenness in moth populations 75–76
- English Channel, Eastern Atlantic (EAT): Eupreptantia 329
- Ensatina* (salamanders): genetics of ice ages 260
- E. eschscholtzii*: molecular data 273 App.
- enzymes
- Bembicium vittatum* (snails): electrophoresis 62
 - conservation strategies, stress and diversity 173–174
 - see also adenosine deaminase and phosphoglucose isomerase
 - development of aquatic insect eggs 232–233
 - Nucella lapillus*: multivariate genetic differentiation 180–181
 - Southeast Asian shrews 200
 - sponge-dwelling alpheid shrimps 310
 - see also loci
- Eocene-Oligocene: first fossil anthomyid 441–442
- Ephemerellidae (Ephemoptera): development of aquatic insect eggs 241 App. 2
- Ephemerella funeralis* 241 App. 2
 - E. subvaria* 241 App. 2
- Ephemeridae (Ephemoptera): development of aquatic insect eggs 241 App. 2
- see also *Hexagenia*
- *Ephemoptera (mayflies): development of aquatic insect eggs 221, 226, 228–233 & Figs 4–5, 240–242 App. 2
 - see also Baetidae, Coloburiscidae, Ecdyonuridae, Ephemerellidae, Ephemeridae, Euthypocidae, Heptageniidae, Leptophlebiidae, Siphonuridae and Tricorythidae
- epilogue: arthropod evolution 47–49
- Epiphyas postvittana*: mtDNA variability and male-lethal symbionts 467
- Epityches eupompe*: ithomiine plant visitors 110 Tab. 5
- equations
- mtDNA variability and male-lethal symbionts 456–459, 462, 470 App.
 - multivariate genetic differentiation 177–180
- equilibrium: mtDNA variability and male-lethal symbionts 453, 455, 461–467 & Figs 4–5 & Tabs 1–2
- Eriobotrya* sp. (Rosaceae): *Aeria olena* visits 106–107 Fig. 2 & Tab. 1
- Ericaceae* (*Vaccinium macrocarpon*): use of AMOVA 360
- Eschweilera*: complementary areas for diversity
- E. ampliplexifolia* 147 App. 2
 - E. baguensis* 149 App. 2
 - E. beebei* 147 App. 2
 - E. boltenii* 149 App. 2
 - E. cabrerana* 149 App. 2
 - E. carinata* 149 App. 2
 - E. compressa* 147 App. 2
 - E. congestiflora* 147 App. 2
 - E. costaricensis* 148 App. 2
 - E. fanshwei* 148 App. 2
 - E. hondurensis* 151 App. 2
 - E. jacquelyniae* 147 App. 2
 - E. klugii* 149 App. 2
 - E. longipedicellata* 147 App. 2
 - E. longirachis* 149 App. 2
 - E. macrocarpa* 147 App. 2
 - E. mexicana* 151 App. 2
 - E. microcalyx* 147 App. 2
 - E. neblinensis* 147 App. 2
 - E. piresii* 148 App. 2
 - E. piauensis* 148 App. 2
 - E. rabeliana* 150 App. 2
 - E. revoluta* 147 App. 2
 - E. roaimensis* 147 App. 2
 - E. rutifolia* 149 App. 2
 - E. subcordata* 148 App. 2
- Esenbeckia leiocarpa* (Rutaceae): pyrrolizidine alkaloids 110 Tab. 5
- esterase (EC 3.1.1.-)
- Bembicium vittatum* (snails): *Est* 4: 62, 65–67 & Fig. 3 & Tab. 2, 72
 - Nucella lapillus* (dogwhelks): *Est*-3 180–181, 183–185 & Tab. 3 & fig. 4
 - Southeast Asian shrews
 - EST (EC 3.1.1.1.) 200
 - Est* 204–206 Tabs 1–3
- Euarthropoda: arthropod evolution 17, 19
- Eueldes* (butterflies): molecular data 274 App.
- Euholognatha: development of aquatic insect eggs 230, 238–239 App. 2
- see also Capniidae, Leuctridae, Nemouridae, Notomouridae and Taeniopterygidae
- Eumalacostraca: compound eyes 41 Fig. 4

- Eupatoriaceae (Asteraceae): pyrrolizidine alkaloids in
 ithomiine butterflies 106 Fig. 2, 117
see also *Ageratum*, *Eupatorium*, *Mikania* and *Trichogonia*
Eupatorium
- flowers: *Apis mellifera* (bees)
E. laevigatum 109
E. maximiliani 109
- pyrrolizidine alkaloids in ithomiine butterflies 100
E. clematideum 109, 113–114 & Tab. 9, 117
E. gaudichaudianum 109, 113–114 & Tab. 9, 117
E. inulaefolium 109, 113–114 & Tab. 9, 117
E. laevigatum 106 Fig. 2, 109–110 & Tab. 5,
 113–114 & Tab. 9, 117
E. maximiliani 106–109 & Figs 2–3 & Tabs 1 & 4,
 113–114 & Tab. 9, 117
E. pauciflorum 106 Fig. 2
E. punctulatum 109, 113–114 & Tab. 9, 117
E. sp. 109 Fig. 3, 114
E. squalidum 109, 113–114 & Tab. 9, 117
- Euphagus cyanocephalus* (Brewer's blackbirds): egg crypsis
 159–162 & Fig. 1, 164–171 & Tabs 1–3 & Figs
 2–4
- Euphorbiaceae: *Aeria olena* visits 108 Tab. 3, 110 Tab.
 5
see also *Croton* and *Dactyloctenion*
- Eupithecia*: evenness in moth populations 76
- Euplectes* (widowbirds): visual signals
E. macrourus (yellow-shouldered) 391–392 Tab. 1
E. prognus (long-tailed) 391–392 Tab. 1
- Eurasia: *Crepis tectorum* 402
- Eureptantia: cladistic biogeography 325–341 & Figs
 1–6 & Tabs 1–2
- Europe: genetics of ice ages 247–266 & Figs 5–7, 275
 App.
see also Alps, Carpathians, Pyrenees and
 Mediterranean Sea
- Eurycerus*: arthropod evolution 3
- Eurythorhynchus pygmaeus*: aggressive nest defence 439
 App.
- Euthypocidae (Ephemoptera): development of aquatic
 insect eggs 241 App. 2
Euthyplocia hecuba 231, 241 App. 2
- evenness (J): moth populations 75–82 & Figs 1 & 3 &
 Tab. 1
- *evolution: cladistic biogeography of the Eureptantia
 325–326, 328
- evolution
 arthropod: reflections 1–55 & Figs 1–4
 growth similarities in tits 343–345, 352, 354
 ornaments are armaments 385–390, 393–394
- *evolutionary constraints: flower-fruit size allometry
 401–402, 406
- Exelodendrum condatum*: complementary areas for
 diversity 148 App. 2
- experiments
 butterfly-ant mutualism 290
 effect of light and diet on a polymorphic caterpillar
 277, 279–280, 283
 ornaments are armaments 390, 392
- Exploring Evolutionary Biology*, by M. Slatkin. Readings
 from *American Scientist*. (Book Review: R. J. Berry)
 500
- *extinction: population genetic effects of a butterfly-
 ant mutualism 287, 289, 300
 extinction/recolonization 287, 301–303
- eyes: arthropod evolution 1–2, 17, 54
 convergence in compound eyes 37–40
 cyclessness in early arthropods 40–44 & Fig. 4
- *F-statistics: multivariate genetic differentiation
 173–176, 179 Tab. 1, 181–184 & Tabs 2–3,
 186–192 & Fig. 6 & Tab. 4
- F-statistics
 analogues: RAPD variation in *Iliamna* 364–365
see also Wright's F-statistics
- Fabaceae (*Crotalaria*): pyrrolizidine alkaloids 100
- Facetotecta: compound eyes 41–42 & Fig. 4
- families: development of aquatic insect eggs 221, 228,
 233
- Fanniidae: Diptera 442
- females
 aggressive nest protection by waders 414, 436–439
 App.
 first fossil anthomyid 441–445 & Figs 1–2
 growth similarities in tits 345–347
 hosts: mtDNA variability and male-lethal symbionts
 453–463 & Fig. 2, 465–465
 daughters 453, 455–456, 460–462
 sisters 455
 ithomiine butterflies: pyrrolizidine alkaloids 99,
 105–108 & Tabs 1–4, 111–112 Tabs 6–8, 118
Jalmenus evagoras (Lycaenidae): butterfly-ant
 mutualism 289–290
 monandry in damselflies 85–93 & Fig. 1 & Tab. 1,
 94–97
 ornaments are armaments 385–394 & Tab. 1
- field techniques: egg crypsis in blackbirds 160–162 &
 Fig. 1
- finches: growth similarities 353–354
 cardueline 352
 fringilid 352–353
 open-nesting 344–345, 353–354
see also chaffinches, greenfinches, rosefinches and
Geospiza
- Finland: genetics of ice ages 252, 258
- Fisher and Ford's Method: survival rates (damselflies)
 92–93
- Fisher, R. A.
 exact tests
 monandry in damselflies 92, 94
 shrimp behavioural responses 312, 318–319 &
 Fig. 5
 run-away models: ornaments are armaments 386
The Genetical Theory of Natural Selection. Conservation
 strategies, stress and diversity 471–472, 475,
 477–479
- fishes
 cladistic biogeography of the Eureptantia *see*
 acanthuroids and gobies
 ornaments are armaments 386, 389–390
see also
 damsel-, pipe- and pupfish, guppies, mollies,
 sticklebacks
Gambusia, *Padogobius* and *Poecilia*
 rainbow (*Melanotaeniidae*): molecular data 276 App.
see also, acanthuroid, sun-, toad- and whitefish
- *fitness: conservation strategies, stress and diversity
 471–473, 475, 477–479
- flatworms: arthropod evolution 25 Fig. 1
- flies

- ornaments are armaments 389
 see also, *Cyrtodictysis*, *Physiophora* and *Scatophaga*
- see also
 black-, butter-, dragon-, fruit-, haw-, may- and stoneflies
 Diptera
- Flora Neotropica*: complementary areas for diversity 125–139 & Figs 1–4 & 6–7
- *floral evolution: flower-fruit size allometry 401
- Flores (Indonesian archipelago): Eupreptantia 327–328 & Fig. 1, 331, 333 Tab. 2, 335 Fig. 4
- Florida: genetics of ice ages 248–250, 257, 262, 272 App, 274 App.
- flower-fruit size allometry: *Crepis* (Asteraceae) 401–407 & Figs 1–2
- Forficula auricularia* (common earwigs): visual signals 391–392 Tab. 1
- *Formicidae: population genetic effects of a butterfly-ant mutualism 287
- Formicoidea: *Nephila* diet 120
- fossils
 arthropod evolution 4–10, 14–15, 21, 23, 29
 computer studies 55
 molecular methods 46–47
 polyphyletic concepts 32–33, 36, 44
 first anthomyid and comments on clade definitions 441–451 & Figs 1–5
- fowl
 domestic (*Gallus gallus*): egg coloration 169
 ornaments are armaments
 domestic 389
 jungle 389
 red jungle (*Gallus gallus*) 391–392 Tab. 1
- foxes see *Vulpia*
- France
 genetics of ice ages 250, 259, 275 App.
 Marseille (Mediterranean, ME): Eupreptantia 329
- Freeman-Tukey method: statistics of synapomorphies 487–489 & Tab. 1, 496–498 App. 2 & Tab. A4
- frogs, Australian *Uperoleia rugosa*: acoustic signals 391–392 Tab. 1
- fruit-flower size allometry: *Crepis* (Asteraceae) 401–407 & Figs 1–2
- fruitflies see *Drosophila*
- FSTAT computer program: multivariate genetic differentiation 181
- Functional Morphology in Vertebrate Palaeontology*, ed. J. J. Thomason (Book Review: Peter Forey) 245–246
- Fundulus heteroclitus* (fundulid fish): molecular data 272 App.
- G-6PD (EC 1.1.1.49): Southeast Asian shrews 200
G-6pd 204–206 Tabs 1–3
- GABAergic innervation of leg muscles: arthropod evolution 35
- Gallinago*: aggressive nest defence
G. gallinago (common snipe) 417 Fig. 2, 438 App.
G. hardwickii 438 App.
G. media 438 App.
G. megal 413, 438 App.
G. nemoricola 413
G. solitaria 413, 438 App.
G. stenura 438 App.
- Gallinago media* (great snipe): visual signals 391–392 Tab. 1
- Gallus gallus*
 domestic fowl: egg coloration 169
 red jungle fowl: visual signals 391–392 Tab. 1
- Gambarellodes* species group: sponge-dwelling alpheid shrimps 308
- Gambusia holbrooki* (fishes): ornaments are armaments 389–390
- Gasterosteus* (sticklebacks): genetics of ice ages 264
G. aculeatus: molecular data 272 App.
- Gastrolina depressa*: mtDNA variability and male-lethal symbionts 467
- Gastropoda
 geographic cohesiveness in an intertidal snail 57, 61, 64 Tab. 1
Nucella lapillus (dogwhelks) 173, 175, 180–182 & Fig. 2, 187–190
- geese: molecular data
Branta canadensis (Canada geese) 273 App.
Chen caerulescens (snow geese) 273 App.
- *gene flow: geographic cohesiveness in an intertidal snail 57–59, 70–72
- gene flow
 genetics of ice ages 248, 256, 259, 266
 specialization and speciation in shrimps 309, 319–320
- genera: development of aquatic insect eggs 221–222, 228
- *genetic bottleneck: mtDNA variability and male-lethal symbionts 453–455, 464, 468
- *genetic structure: effects of a butterfly-ant mutualism 287–290, 292–295 & Tabs 2–3, 303
- genetics
 developmental: arthropod evolution 26–30, 33, 40
Distal-less (Dll) 7, 29–30
 differentiation: SE Asian shrews 197–199, 201–208 & Tabs 1–3, 212–215 & Fig. 5
 see also distance-Wagner, Nei's and Rogers' procedures
 diversity: conservation strategies and stress 471–482
 ice ages 247–248, 251–260 & Figs 1–4, 262–263
 molecular data: *Heliconius erato* 274 App.
 population: multivariate genetic differentiation 173–192 & Figs 2–3 & 5–6 & Tabs 1 & 4
 gene flow 174–175, 181, 188–189, 191
 variations: geographic cohesiveness in an intertidal snail 57–59, 62–63, 71–73
- genomes: ice ages 247, 253–256, 259–266 & Figs 6–7
- genus: first fossil anthomyid 441, 447–450
- *geographic cohesiveness: intertidal snails (*Bembicium vittatum*) 57–58
 patterns 62
- geographic distances: SE Asian shrews 197, 202
- *Geometridae: polymorphic caterpillars
 loopers 278
Nemoria arizonaria (emerald moths) 277
- Geospiza* finches: growth 352–353
- gibbons (*Hylobates agilis*): acoustic signals 391–392 Tab. 1
- gill disease: *Onchorhynchus mykiss* (rainbow trout) 474
- Glareola*: aggressive nest defence
G. lactea 413
G. maldiviarum 436 App.
G. nordmanni 436 App.
G. pratensis 436 App.
- Glomeris* (pill millipedes): convergence 35

- Glossolepis*: molecular data 276 App.
glutamate oxaloacetate transaminase (EC 2.6.1.1)
Bembicium vittatum (snails)
Got 1: 62, 65, 72
Got 2: 62
Southeast Asian shrews
Got-1 204 Tab. 1
GOT-1: 200
GOT-2: 200, 203
gobies, sicydine: relation ships between modern oceans 340
Gochnatia polymorpha (Mutisieae, Asteraceae): visits by Ithomiinae 109, 110 Tab. 5
godwhits: aggressive nest defence 427
black-tailed (*Limosa limosa*) 411, 437 App.
Gondwana breakup: eurentatic phylogenetic trees 339
gophers, pocket *see* *Thomomys*
GPD (EC 1.1.1.8): Southeast Asian shrews 200
Gpd 204–205 Tabs 1–2
Gpi (EC 5.3.1.9): butterfly-ant mutualism 291, 295
Tabs 2–3, 299 Tab. 5
*graphical display: multivariate genetic differentiation 173
grass
cheatgrass: spread in North America 251
North American barnyard (*Echinochloa microstachya*): genetics of ice ages 260
grasshoppers, meadow *see* *Chorthippus parallelus*
grassland: evenness in moth populations 75–76, 80–82 & Tab. 1
Great Barrier Reef (Gbr), Australia: Eurentatic 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
*great tits (*Parus major*): growth similarities 343–354 & Tabs 1–5 & Fig. 1
Greece: genetics of ice ages 251–252, 258–259, 263–264
greenfinches
open-nesting 344
ornaments are armaments 387
Greenland: Sirius Passet fauna 15
Greya (moths)
molecular data 274 App.
switch of umbellifer hosts
G. solenobiella 266
G. suffusa 266
Grias multinerva: complementary areas for diversity 148 App. 2
groundplans: arthropod evolution 1, 7–10
bauplane 23, 27
groups: monandry in damselflies 85, 95
Group I 88–97 & Tab. 1
see also *I. Ischnura aurora*, *I. demorsa*, *I. hastatum*, *I. kellicotti*, *I. perparva*, *I. posita* and *I. verticalis*
Group II 88–96 & Tab. 1
see also *I. cervula*, *I. damula*, *I. elegans*, *I. graellsii*, *I. erratica*, *I. pumilio*, *I. ramburi* and *I. senegalensis*
Group III 88, 90–93 & Tab. 1, 95
see also *I. denticollis* and *I. gemina*
grouse
red: ornaments are armaments 387
willow (*Lagopus lagopus*): aggressive nest defence 429
*growth: similarities in tits 343–346, 349–354 & Tabs 3–4 & Fig. 1
Guatemala: genetics of ice ages 249–250
gulls: nest predators of waders 416, 422, 427, 436–439 App.
guppies (*Poecilia reticulata*)
molecular data 272 App.
visual signals 391–392 Tab. 1
gurami, croaking (*Trichopsis vittatus*): acoustic signals 391–392 Tab. 1
Gustavia: complementary areas for diversity
G. acuminata 147 App. 2
G. acuta 151 App. 2
G. coriacea 150 App. 2
G. erythrocarpa 150 App. 2
G. excelsa 149 App. 2
G. f. var. costata 149 App. 2
G. f. var. flagellata 149 App. 2
G. fosteri 147 App. 2
G. gigantophylla 149 App. 2
G. gracillima 150 App. 2
G. inakua 149 App. 2
G. latifolia 150 App. 2
G. longepetiolata 150 App. 2
G. longifuniculata 150 App. 2
G. m. paucisperma 150 App. 2
G. monocaulis 149 App. 2
G. parviflora 149 App. 2
G. petiolata 149 App. 2
G. pubescens 148 App. 2
G. s. speciosa 149 App. 2
G. serrata 149 App. 2
G. sessilis 148 App. 2
G. tejerae 150 App. 2
G. terminaliflora 149 App. 2
Habelia: Cambrian arthropods 23
habitats: geographic cohesiveness in an intertidal snail 57–65 & Figs 1–2 & Tab. 1, 67–68, 70–71, 73
Haematopus: aggressive nest defence
H. bachmani 436 App.
H. ostralegus 436 App.
H. palliatus 436 App.
Haematothemia: statistics of synapomorphies 489, 492
Haplopoda: arthropod evolution 11
haplotypes: genetics of ice ages 247, 258–259
Hardy-Weinberg equilibrium: multivariate genetic differentiation 191
Harmonia axyridis: mtDNA variability and male-lethal symbionts 467
Harpacticoida: arthropod evolution 4, 13
Hartley's *F* test: development of aquatic insect eggs 228
Hawaii: molecular data
snow geese 273 App.
spiders 273 App.
hawflies (*Rhagoletis pomonella*): genetics of ice ages 266
hawkmoths (Sphingidae): colour polymorphism 278
hazel pollen cores: Europe 250
HBB, HK-1 and HK-2 (EC 2.7.1.1): SE Asian shrews 200
Hbb 204 Tab. 1
Hk-1 and *Hk-2* 204–206 Tabs 1–3
Hecatesia thyridion (whistling moths): acoustic signals 391–392 Tab. 1
Heinrich events: genetics of ice ages 249

- Heliantheae (Asteraceae): pyrrolizidine alkaloids 106
 Fig. 2, 114
Emilia sonchifolia 106 Fig. 2, 114
- Heliconiini: pyrrolizidine alkaloids 109, 119–120
- Heliconius* (Lepidoptera)
 convergence
H. erato and *H. melpomene* 35–36
H species 36
 genetics of ice ages 262, 264
 molecular data 274 App.
- heliospathine: pyrrolizidine alkaloid 114
- Heliothis* (moths): mtDNA variability and male-lethal symbionts 467
- Heliotropium* (Boraginaceae): pyrrolizidine alkaloids 100, 114
H. indicum 109 Fig. 3, 113–114 & Tab. 9
H. peruvianum 108 & Fig. 3, 210 Tab. 5, 113 Tab. 9
H. spathulatum 114
H. transversum 105–110 & Fig. 2 & Tabs 1 & 3 & 5, 113–114 & Tab. 9, 117–118
- helminth parasites: stingrays 340
- Helobdella* (leeches): engrailed class genes 29
- Hennigian analyses
 arthropod evolution 48
 statistics of synapomorphies 483–484, 489, 491, 493
- Heptageniidae: development of aquatic insect eggs
 241 App. 2
see also Rhithrogena
- Hesslandona*: Cambrian arthropod 26, 41 Fig. 4
- heterophylline: *Tithorea harmonia* larvae and adults 110
- Heteroscelus*: aggressive nest defence
H. brevipes 438 App.
H. incanus 438 App.
- *heterozygote advantage: conservation strategies, stress and diversity 471–475, 477
- Hexagenia* (Ephemeridae): development of aquatic insect eggs
H. bilineata 240 App. 2
H. rigida 240 App. 2
- Hexapoda: arthropod evolution 2, 19, 34 Fig. 3, 43
- Himantopus* aggressive nest defence
H. himantopus (black-winged stilts) 417 Fig. 2, 436 App.
H. mexicanus 436 App.
- Hirtella*: complementary areas for diversity
H. adderleyi 149 App. 2
H. adenophora 151 App. 2
H. angustissima 148 App. 2
H. araguanensis 147 App. 2
H. aramangensis 150 App. 2
H. bamebyi 148 App. 2
H. barnesoi 147 App. 2
H. confertiflora 147 App. 2
H. cordifolia 147 App. 2
H. corymbosa 147 App. 2
H. couepiflora 149 App. 2
H. cowanii 148 App. 2
H. dorvalii 150 App. 2
H. enneandra 148 App. 2
H. floribunda 150 App. 2
H. guatemalensis 149 App. 2
H. juruensis 150 App. 2
H. kuhlmannii 151 App. 2
H. lancifolia 148 App. 2
H. liesneri 149 App. 2
H. longifolia 149 App. 2
H. magnifolia 148 App. 2
H. maguirei 149 App. 2
H. orbicularis 149 App. 2
H. papillata 147 App. 2
H. parvunguis 147 App. 2
H. pendula 148 App. 2
H. radamii 148 App. 2
H. rugosa 150 App. 2
H. santosii 150 App. 2
H. scaberula 149 App. 2
H. standleyi 150 App. 2
H. subglanduligera 150 App. 2
H. trichotoma 148 App. 2
H. vesiculosa 150 App. 2
- Holarctic (*Osmerus*: smelt): molecular data 273 App.
- homeobox genes: arthropod evolution 26–27
- homologies: arthropod evolution 1, 3, 36, 39
- HOMOVA (homogeneity of molecular variance analysis): RAPD variation in *Iliamna* 357, 360–361, 365
- homozygosity
 conservation strategies, stress and diversity 472, 474, 477
 genetics of ice ages 247, 253–254
- honey-bees: ornaments are armaments 388
- *Hoploneuracea: cladistic biogeography of the Eurentania 325
- hosts, sponge: specialization and speciation in shrimps 307–312 & Fig. 1, 314–321 & Figs 4–5
- *hotspots: complementary areas for diversity 126–128
- Houtman Abrolhos Islands, Western Australia:
Bembicium vittatum (snails) 57–60 & Fig. 1, 66 Fig. 3, 70–72
see also Easter, Pelsaert and Wallabi Groups 60 Fig. 1
- Hox genes: arthropod evolution 33–34 & Fig. 3
- Hubrechtionemertes lanksteri*: eurentantic nemertean (WAT) 327 Fig. 1, 329, 332 Fig. 2, 339
- Hyalurga syma* (Arctiidae: moths): pyrrolizidine alkaloids 114, 118
- Hyatella intestinalis*: host sponge 315–316
- Hybanthus atropurpureus* (Violaceae): *Aeria olenae* visits 106–107 Fig. 2 & Tab. 1
- *hybrid zones: genetics of ice ages 247–248, 256–261 & Figs 5–6, 263–266 & Fig. 7
- hydroids: hermit crab shell crusts 321
- Hydrophasianus chirurgus*: aggressive nest defence 413
- Hylobates agilis* (gibbons): acoustic signals 391–392 Tab. 1
- Hymeniacidon* cf. *caerulea*: host sponge 315, 318 Fig. 4
- Hymenoptera
Heliotropium transversum flowers 109
Nephila diet 120
- hypotheses
 aggressive nest defence: alternative prey 409, 411–412, 417, 423–426 & Tabs 1–2, 428–429
 arthropod evolution 10, 13–14, 18–19, 22
 cyclessness in early arthropods 40–44 & Fig. 4
 cladistic biogeography of the Eurentania 325–328 & Fig. 1, 329–331, 335, 338–341
 development of aquatic insect eggs 226 Fig. 2
 egg crypsis in blackbirds 159–161 & Fig. 1
 first fossil anthomyid and comments on clade definitions 447

- flower-fruit size allometry 402, 405–406
ornaments are armaments 393
pruinescence and predation (damselflies) 95–96
SE Asian shrews 197, 199, 210, 212 Figs 5–6, 214
 see also Brook's parsimony analysis
specialization and speciation in shrimps 307, 321
statistics of synapomorphies 484–486 & Figs 1–2, 492–493
 see also null hypothesis
Hypothyris ninomia dasta: ithomiine plant visitors 110
 Tab. 5
hyrax, rock (*Procavia capensis*): molecular data 275 App.
- Iberia: genetics of ice ages 252, 258, 263–265 & Fig. 7
Ibidorhyncha struthersii: aggressive nest defence 436 App.
ice ages: genetic consequences 247–251, 258–266 & Figs 5 & 7
Icteridae (blackbirds): egg crypsis 159–162 & Fig. 1
 see also *Agelaius*, *Euphagus* and *Xanthocephalus*
IDH-1 and IDH-2 (EC 1.1.1.42)
 SE Asian shrews 200
 Idh-1 and *Idh-2* 205 Tab. 2
 IDH-2 204 Tab. 1, 206 Tab. 3
sponge-dwelling alpheid shrimps
 Idh 310, 312 Fig. 1
 Idh-1 313 Tab. 1
Iliamna (mallows): RAPD variation 357–361
 I. corei (Peters Mountain) 357–365 & Figs 1–3 & Tabs 1–3
 I. remota var. *corei* 359
 Phymosia remota 358–359
 I. remota (Kankakee) 357–365 & Figs 1–3 & Tabs 1–3
 I. remota var. *typica* 359
 I. rivularis 360
Ilione albiseta (Sciomyzidae): development of aquatic insect eggs 228, 242 App. 2
Illinois: *Ischnura kellycotti* 89
Inaequifurcata: division of the eurentic nemertean 326, 338
inbreeding depression: conservation strategies, stress and diversity 472–473
incubation: aggressive nest protection 427–428, 326–329 App.
 sharing 409, 420 Fig. 4, 427
 times 409–411, 414–415, 418–420, 424–425 & Tab. 1, 429, 436–439 App.
India: *Suncus dayi* (shrews) 199
Indian Ocean: cladistic biogeography of the Eurentia 328, 330–331
 Eastern (EIO) 325, 327–340 & Figs 1–6 & Tab. 1
 see also Indonesian archipelago: *Coella*, *Drepanobanda*, *Drepanophorella*, *Drepanophoria*, *Drepanophorina*, *Drepanophoringia*, *Drepanophorus*, *Paradrepanophorus*, *Punnettia* and *Sibogamermes*
 Western (WIO) 325, 327–328 & Fig. 1, 330–340 & Tab. 1 & Figs 1–6
 see also Maldives: *Drepanophorella* and *Punnettia*
indicator models: ornaments are armaments 386, 388
indices
 cladistic biogeography of the Eurentia
 consistency (CI) 331–332, 334–338 & Figs 3–6 & Tab. 3
 retention (RI) 331–332, 334–337 Figs 3–6 & Tab. 3
evenness in moth populations 79–80
 see also Berger-Parker, Shannon and Simpson
exposure: geographic cohesiveness in *Bembicium vittatum* (snails) 61
retention (chromatography): pyrrolizidine alkaloids in ithomiine butterflies 105
indicine (pyrrolizidine alkaloid): ithomiine butterflies 99, 102 Fig. 1, 114, 117
Indo-Pacific: cladistic biogeography of the Eurentia 327, 331–332, 339–340
*Indomalayan region: SE Asian shrews 197, 199, 213
Indonesian archipelago: cladistic biogeography of the Eurentia 328, 338–341
 see also Ambon, Flores, Obi, Savu, Seram, Sulawesi, Sumba, Timor and Tiur
infection: mtDNA variability and male-lethal symbionts 453–467 & Figs 1–2 & 5
Insecta
 aquatic insects
 development of eggs 221–244 & Figs 1–5 & Apps 1–2
 monandry in damselflies 85
arthropod evolution 8, 16, 18, 21, 38
 faceted eyes 39
 groundplan of insect/arthropod leg 1, 7–10
genetics of ice ages 251, 258–259 & Fig. 5, 262, 266
 see also blackflies, butterflies, *Chorthippus* and hawflies
mtDNA variability and male-lethal symbionts 454–455, 459, 467–468
ornaments are armaments 386
 see also cockroaches, crickets, earwigs, flies, honeybees, moths, spiders and swordtails
phytophagous and zooparasitic insects: speciation 321
Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA): complementary areas for diversity 127, 129
Instituto Nacional de Pesquisas de Amazonia (INPA): complementary areas for diversity 127, 129, 137
integerrimine (pyrrolizidine alkaloid): ithomiine butterflies 111
*intermediate (pyrrolizidine alkaloid): ithomiine butterflies 99–101, 102 Fig. 1, 104, 109–110, 114, 117
Internet: multivariate genetic differentiation 192
Interstate 64(I): RAPD variation in *Iliamna* 358 Fig. 1, 360, 362–365 & Tabs 1–3 & Fig. 2
introductions
 aggressive nest protection by waders 410–412
arthropod evolution 1–2
cladistic biogeography of the Eurentia 325–326
complementary areas for diversity 126–128
conservation strategies, stress and diversity 471–473
development of aquatic insect eggs 222–223
effect of light and diet on a polymorphic caterpillar 277–279
egg crypsis in blackbirds 160
evenness in moth populations 75–76
first fossil anthomyid and comments on clade definitions 442
flower-fruit size allometry 401–402
genetics of ice ages 48
geographic cohesiveness in an intertidal snail 58–59

- growth similarities in tits 343–345
 monandry in damselflies 85–86
 mtDNA variability and male-lethal symbionts 454–455
 multivariate genetic differentiation 174–176
 ornaments are armaments 386
 population genetic effects of a butterfly-ant mutualism 288–290
 pyrrolizidine alkaloids in ithomiine butterflies 100–101
 RAPD variation in *Iliamna* 358–360
 SE Asian shrews 198–199
 specialization and speciation in shrimps 308–309
 statistics of synapomorphies 483–484
 introgression: genetics of ice ages 248, 258–269
 invertebrates, marine: specialization and speciation in shrimps 307–308, 320
 Ireland: genetics of ice ages 251, 275 App.
 **Iridomyrmex*: population genetic effects of a butterfly-ant mutualism 287, 292 & Tab. 1, 300, 303
I. anceps 288–292 & Fig. 1 & Tab. 1, 298–301 & Fig. 4
I. ricinus 288–292 & Fig. 1 & Tab. 1, 298–299 & Fig. 4
 Iron Gate (IG), Virginia: RAPD variation in *Iliamna* 358 Fig. 1, 360, 362–363 Tabs 1–2, 365 & Tab. 3
Ischnura: monandry in damselflies 85–97 & Tab. 1
I. aurora 87–88, 90–91 & Fig. 1 & Tab. 1, 93–96
I. cerulea 88, 90–92 & Fig. 1 & Tab. 1, 94
I. damula 88, 90–92 & Fig. 1 & Tab. 1, 94
I. demorsa 87–88, 90–91 & Fig. 1 & Tab. 1, 93–94, 96
I. denticollis 88, 90–91 & Fig. 1 & Tab. 1, 93
I. elegans 88, 90–93 & Fig. 1 & Tab. 1
I. erratica 88, 90–91 & Fig. 1 & Tab. 1
I. gemina 88, 90–91 Fig. 1 & Tab. 1, 93
I. graellsii 87–89, 90–93 & Fig. 1 & Tab. 1
I. hastatum: Citrine Forktail 87–88, 90–94 & Fig. 1 & Tab. 1
I. kellicotti 87–89, 90–92 & Fig. 1 & Tab. 1, 94
I. perparva 87–88, 90–91 & Fig. 1 & Tab. 1, 94
I. posita 87–88, 90–94 & Fig. 1 & Tab. 1
I. pumilio 88, 90–93 & Fig. 1 & Tab. 1
I. ramburi 88, 90–91 & Fig. 1 & Tab. 1
I. senegalensis 88, 90–91 Fig. 1 & Tab. 1
I. verticalis 87–88, 90–94 & Fig. 1 & Tab. 1, 96
Ischnura verticalis (Coenagrionidae): development of aquatic insect eggs 240 App. 2
 island groups: geographic cohesiveness in an intertidal snail 57–65 & Figs 1–2, 67–72 & Tabs 2–4 & Figs 4–5
 isocallimorphine: pyrrolizidine alkaloid 117
Isogenus nubecula (Periodicidae): development of aquatic insect eggs 230, 239 App. 2
 *isolation by distance: geographic cohesiveness in an intertidal snail 57, 70
 isometry: flower-fruit size allometry 401–405
Isonychia bicolor (Siphonuridae): development of aquatic insect eggs 241 App. 2
 isopelletierine (pyrrolizidine alkaloid): ithomiine butterflies 111
Isoperla (Periodicidae): development of aquatic insect eggs
I. diffinis 228, 239 App. 2
I. grammica 228, 230, 239 App. 2
 Israel: *Spalax ehrenbergi* (mole rats) 275 App.
 Italy
 genetics of ice ages 250–251, 258–259, 264–265 & Fig. 7, 275 App.
 Mediterranean (ME): Eureptantia 329
Ithomia agnosia: Ithomiinae plant visitors 110 Tab. 5
 Ithomiinae (butterflies): pyrrolizidine alkaloids 99–123 & Figs 1–3 & Tabs 1–11
 see also *Aeria*, *Ithomia*, *Mechnitis* and *Tithorea*
Jacana spinosa: aggressive nest defence 413
 jackknife procedures (analysis): butterfly-ant mutualism 293, 295 Tabs 2–3
Jalmenus (Lepidoptera: Lycaenidae): butterfly-ant mutualism 288
J. daemelii 288
J. exagoras 287–295 & Fig. 1 & Tabs 2–3, 297–303 & Tab. 5
J. icinus 288
 Java: SE Asian shrews 197–198, 200 Fig. 1, 202–203 & Fig. 2, 205 Tab. 2, 208–212 & Figs 5–6, 214–215, 219 App.
 jaws: arthropod evolution 1–7
 Jolly-Seber Stochastic Method: survivorship (damselflies) 92
Jonidium brevicaule (Violaceae): *Aeria olenae* visits 106–107 Fig. 2 & Tab. 1
 junco, dark-eyed: ornaments are armaments 387
 Jurassic
 arthropod evolution 33
 eureptantian phylogenetic trees 339
Kerygmachela: arthropod evolution 12, 15
K. kærkegaardi 15
Kobus vardonii (Puku antelope): visual signals 391–392 Tab. 1
Kohlrusteria paniculata (Sapindaceae): visits by Ithomiinae 109–110 & Tab. 10
 Kruskal-Wallis H: monandry in damselflies 89, 93
 *labrum: arthropod evolution 1, 7, 16–19
 Lactuceae, Asteraceae (*Crepis*): flower-fruit size allometry 402
 ladybirds, two-spot (*Adalia bipunctata*): cytoplasmic male-killers 459, 462–463, 467–468
Lagopus
L. lagopus (willow grouse): aggressive nest defence 429
L. mutus (rock ptarmigans): visual signals 391–392 Tab. 1
 Lambert, D. M. and Spencer, H. G. *Speciation and the Recognition Concept, Theory and Application*. (Book Review: M. F. Claridge) 499
 Lannacombe Bay (Devon coast): *Nucella lapillus* (dogwhelks) 181–183 & Figs 2–3, 185–187 & Figs 5–6
Lantana sp. (Verbenaceae): *Aeria olenae* visits 106–107 Fig. 2 & Tab. 1
Lap-1 & *Lap-2*, allozyme variations (*Nucella lapillus*: dogwhelks) 180–181, 183 Tab. 3, 185 Fig. 4
Lap-2 183
 lapwings: aggressive nest defence 422, 427
Vanellus vanellus 411, 417 Fig. 2, 436 App.
 larvae
 development of aquatic insect eggs 223, 230–233

- I. obscura* 228, 230, 232, 239 App. 2
 effect of light and diet on a polymorphic caterpillar:
Nemoria arizonaria 277–281, 283
 population genetic effects of a butterfly-ant
 mutualism 288–290, 300
 pyrrolizidine alkaloids in ithomiine butterflies
 99–101, 103–104, 109–112 & Tabs 6–8, 114,
 118
 latitudes (breeding): aggressive nest protection by
 waders 409–412, 415, 420–423–425 & Fig. 6 &
 Tab. 1, 429
 Laurentide ice sheet: genetics of ice ages 249–250, 261
 LDH-1 and LDH-2 (EC 1.1.1.27.): SE Asian shrews
 200, 203
 leaves, oak: effect of light and diet on a polymorphic
 caterpillar 277, 279–283 & Tab. 1 & Fig. 2
 Lecythidaceae: complementary areas for diversity 125,
 128–129, 132–138 & Figs 1 & 3–4 & 6–7
Lecythis aurantiflora 148 App. 2
L. mesophylla 149 App. 2
L. pneumatophora 147 App. 2
L. schwackei 151 App. 2
L. serrata 151 App. 2
 leeches (*Helobdella*): engrailed class genes 29
 legs, insect/arthropod: groundplan 7–10
 Leguminosae: pyrrolizidine alkaloids in ithomiine
 butterflies 106–108 Fig. 2 & Tabs 1 & 3, 110
 Tab. 5, 118
 see also *Acacia*, *Calopogonium* and *Crotalaria*
Leoneohtilia: Cambrian arthropods 23, 26
Lepidocaris: Devonian lipostracan 3, 55
 *Lepidoptera: evenness in moth populations 75–84 &
 Figs 1–3 & Tab. 1
 Lepidoptera
 arthropod evolution
Heliconius erato and *H. melponome* 35–36
Precis coenia 7
 butterfly-ant mutualism (*Talmenus evagoras*:
Lycanidae) 287
 mtDNA variability and male-lethal symbionts 468
Nemoria arizonaria (emerald moths) 277–278, 283
 pyrrolizidine alkaloids in ithomiine butterflies
 99–100, 118, 120
 see also Arctiidae (moths), danaine and ithomiine
 butterflies
 see also Macrolepidoptera and moths
 Lepidosauria: statistics of synapomorphies 490–492 &
 Tabs 2–3
Lepidurus: arthropod evolution 3
Lepomis (sunfish): genetics of ice ages 264
L. gulosus (Warmouth) 272 App.
L. macrochirus (bluegill): molecular data 272 App.
L. microlophus (redeer) 272 App.
L. punctatus (spotted) 272 App.
Leptodora: eye structure 38
 Leptophlebiidae (Ephemeroptera): development of
 aquatic insect eggs 241 App. 2
 see also *Nousia*
Lestes (Odonata): development of aquatic insect eggs
 233
 leucine aminopeptidase (EC 3.4.–)–
Bembicium vittatum (snails), *Lap-1* 62, 65, 67–68 &
 Tab. 2, 72
Mytilus edulis: *Lap* polymorphism 58
Leucorrhina (Libellulidae): development of aquatic
 insect eggs
L. borealis 240 App. 2
L. glacialis 240 App. 2
L. proxima 240 App. 2
 Leuctridae (Plecoptera): development of aquatic insect
 eggs 238 App. 2
Leuctra digitata 238 App. 2
L. fusca 238 App. 2
L. geniculatus 238 App. 2
L. hippopus 228, 238 App. 2
L. inermis 238 App. 2
L. moselyi 238 App. 2
L. nigra 238 App. 2
 leucylglycylglycine peptidase (EC 3.4.1.3): *Bembicium*
vittatum (snails) *Lapp* 62, 67 & Tab. 2
 leucylproline peptidase (EC 3.4.1.3): *Bembicium vittatum*
 (snails) *Lpp* 62, 72
 Libellulidae (Odonata): development of aquatic insect
 eggs 240 App. 2
Libellula julia 240 App. 2
 see also *Leucorrhina*
Licania: complementary areas for diversity
L. angustata 150 App. 2
L. anneae 150 App. 2
L. apiculata 150 App. 2
L. aracaensis 148 App. 2
L. araneosa 148 App. 2
L. arianeae 148 App. 2
L. bahiensis 150 App. 2
L. bellingtonii 148 App. 2
L. boliviensis 151 App. 2
L. buxifolia 148 App. 2
L. cabrae 148 App. 2
L. cardiophylla 148 App. 2
L. cari 147 App. 2
L. cecidiophora 150 App. 2
L. chiriquiensis 149 App. 2
L. cidii 148 App. 2
L. compacta 147 App. 2
L. conferruminata 150 App. 2
L. couepizfolia 150 App. 2
L. cyathodes 147 App. 2
L. cymosa 148 App. 2
L. dealbata 148 App. 2
L. dodsonii 151 App. 2
L. durifolia 148 App. 2
L. espiniae 150 App. 2
L. filomenoi 150 App. 2
L. foldatsii 148 App. 2
L. foveolata 147 App. 2
L. fuchsii 149 App. 2
L. fusfuracea 147 App. 2
L. gentryi 150 App. 2
L. glazioviana 147 App. 2
L. gonzalezii 151 App. 2
L. guatemalensis 149 App. 2
L. hebantha 149 App. 2
L. hedbergii 149 App. 2
L. hispida 150 App. 2
L. hitchcockii 150 App. 2
L. hypoleuca 132–133 Fig. 2
L. imbamadaensis 147 App. 2
L. jaramilloi 149 App. 2
L. joseramosii 150 App. 2

- L. krukovi* 148 App. 2
L. latistipula 148 App. 2
L. maguirei 150 App. 2
L. maranhensis 150 App. 2
L. maritima 149 App. 2
L. maxima 147 App. 2
L. megalophylla 149 App. 2
L. mexicana 150 App. 2
L. michauxii 151 App. 2
L. microphylla 151 App. 2
L. miltonii 150 App. 2
L. minuscula 149 App. 2
L. montana 150 App. 2
L. morii 147 App. 2
L. nelsonii 148 App. 2
L. niloi 148 App. 2
L. operculipetala 147 App. 2
L. pakaraimensis 149 App. 2
L. pittieri 147 App. 2
L. pruinosa 147 App. 2
L. retifolia 149 App. 2
L. rigida 150 App. 2
L. voraimensis 147 App. 2
L. sandwithii 148 App. 2
L. santosii 147 App. 2
L. silvatica 149 App. 2
L. stevensii 148 App. 2
L. stricta 150 App. 2
L. subrotundata 147 App. 2
L. tachirensis 149 App. 2
L. tambopatensis 151 App. 2
L. texeirae 150 App. 2
L. tepuiensis 147 App. 2
L. ternatensis 148 App. 2
L. tocanina 149 App. 2
L. vasquezii 150 App. 2
L. vaupesiana 151 App. 2
L. velutina 150 App. 2
- lice (parasites): feral rock doves 474
- light effects: *Nemoria arizonaria* (emerald moths)
 277 283 & Fig. 1 & Tab. 1
- *limbs: arthropod evolution 1–10
 groundplan 7–10
 primitive arthropods 10–15
- Limicola falcinellus*: aggressive nest defence 439 App.
- Limnia unguicornis* (Sciomyzidae): development of
 aquatic insect eggs 242 App. 2
- Limnocyrtus minimus*: aggressive nest defence 438 App.
- Limnodromus*: aggressive nest defence
L. griseus 438 App.
L. scolopaceus 438 App.
L. semipalmatus 438 App.
- Limoniinae (Diptera): development of aquatic insect
 eggs 231
- Limosa*: aggressive nest protection
L. fedoa 437 App.
L. haemastica 437 App.
L. lapponica 437 App.
L. limosa (black-tailed godwit) 411, 437 App.
- limpets, pulmonate (*Siphonaria kurracheensis*): intertidal
 communities 61, 64 Tab. 1
- Limulus*
 compound eyes 38, 42
L. polyphemus (horseshoe crabs): molecular data 272
 App.
- *lineage: first fossil anthomyid and comments on clade
 definitions 441, 448
- Linyphia litigiosa* (spiders): ornaments are armaments
 390
- Lissamphibia: statistics of synapomorphies 490–492 &
 Tabs 2–3
- Lissodendoryx* (sponges)
L. cf. strongylata: Belize 309–310, 312–314 & Fig. 1,
 316, 318 Fig. 4, 321
L. colombiensis: Panama 309–315 & Fig. 1, 317–319
 & Figs 4–5
- Litorina*: localised selection
L. mariae 58
L. saxatilis 58
- Littoraria undulata*: geographic cohesiveness in intertidal
 snails 61, 63 64 & Tab. 1
- *littorine snails: geographic cohesiveness in *Bembicium*
vittatum 57–58, 61
- lizards
Anolis carolinensis: visual signals 391–392 Tab. 1
Sceloporus occidentalis: malarial parasitic stress 474
- loci, enzyme
 butterfly-ant mutualism 290–291, 295 Tabs 2–3,
 299 & Tab. 5
see also Aat-1, AcoH-1, Ak, Gpi, Pgm-1 and Sod
 genetics of ice ages 253, 255 Fig. 2, 258, 264–265
 & Fig. 7
 geographic cohesiveness in an intertidal snail 57–59,
 62–63, 65–66 & Fig. 3, 71–72
 multivariate genetic differentiation 173–185 & Figs
 1 & 4 & Tab. 3, 187, 189–192 & Tab. 4
 SE Asian shrews 197, 200–201, 203–206 & Tabs
 1–3, 213
see also ADA, ADH, AK, ALB, CK, EST, GOT,
GPD, G-6PD, HBB, HK, IDH, LDH, MDH,
MOH, MPI, PA, 6-PGD, PGI, PGM, PROT,
SOD and XDH
 sponge-dwelling alpheid shrimps 310, 312–314 &
 Fig. 1 & Tab. 1, 319–320
see also Aat, Idh, Mdh-1, Mdh-2, Mpi, Pgi, Pgm and
Tpi
 stress 474–476
- log-transformations
 evenness in moth populations 75, 77, 79–81
 growth similarities in tits 345–346, 349 Tab. 3
- loopers (Geometridae): polymorphism of caterpillars
 278
- Lucilia cuprina* (insects): intraspecific competition 478
- Lycanidae (*Jalmenus evagoras*: Lepidoptera): butterfly-
 ant mutualism 287–290, 303
- *lycopsamine (pyrrolizidine alkaloid): ithomiine
 butterflies 99–102 & Fig. 1, 104, 109–110,
 113–114, 117
- Macaca* (macaques): molecular data
M. cyclopsis 276 App.
M. fuscata 276 App.
M. mulatta 276 App.
M. sinica group 276 App.
- MacClade 3.0 (computer program): cladistic
 biogeography of the Eureptantia 328, 333
- Macrolepidoptera: evenness in moth populations
 75–84 & Figs 1–3 & Tab. 1
- Madracis mirabilis*: reef corals 309
- Magicicada* (periodic cicadas): molecular data 274 App.

- *major axis regression: flower-fruit size allometry 401–402, 404
- Malaclemys*: genetics of ice ages 262
- M. terrapin* ("Diamondback terrapin"): molecular data 272 App.
- Malacostraca: arthropod evolution 7–8, 16, 18, 28–29, 41
- Myxidopsis* 7
- malarial parasites (*Plasmodium mexicanum*): lizards 174
- malate dehydrogenase pattern: *Culex pipiens fasciatus* 232
- Malay Archipelago: SE Asian shrews 197–203 & Figs 1–2, 210–211, 213–216
 - peninsula 197–198, 200 Fig. 1, 202–204 & Fig. 2 & Tab. 1, 208–212 & Figs 5–6, 215, 219 App.
- Maldives, West Indian Ocean (WIO): Eupreptantia 327–328 & Fig. 1, 331, 333 Tab. 2, 335 Fig. 4
- Malenka californica* (Nemouridae): development of aquatic insect eggs 238 App. 2
- males
 - aggressive nest protection by waders 414, 436–439 App.
 - butterfly-ant mutualism: *Jalmenus evagoras* (Lycaenidae) 289–290
 - growth similarities in tits 345–347
 - ithomiine butterflies: pyrrolizidine alkaloids 99–100, 105–108 & Tabs 1–4, 111–112 & Tabs 6–8, 114, 117–118, 120
 - male-lethal symbionts: mtDNA variability 453–468 & Figs 1–6 & Tabs 1–3 & App.
 - monandry in damselflies 85–97 & Fig. 1 & Tab. 1
 - ornaments are armaments 385–394 & Tab. 1
 - male-male combat/competition 385–386, 388–390, 393–394
- *mallow (*Iliamna*): RAPD variation 357–369 & Figs 1–4 & Tabs 1–3
 - Kankakee *see I. remota*
 - Peters mountain *see I. corei*
- Malvaceae: RAPD variation in *Iliamna* 357–358
- mammals
 - aggressive nest protection by waders 413
 - ornaments are armaments 387
 - see also* gibbons, seals and sheep
 - statistics of synapomorphies 489–492
- manakins, swallow-tailed (*Chiroxiphia linearis*): visual signals 391–392 Tab. 1
- Manaus, Brazil: complementary areas for diversity 125, 134, 137
 - E. and W. 144–145 App. 1, 148 App. 2, 152 App. 3, 155 App. 4
- *mandibles: arthropod evolution 1, 17–20, 22, 39, 48 nature 2–7
 - origin 10–15
 - protomandibles 19–20
- Mandibulata: arthropod evolution 1–7, 13, 17, 19–20, 39, 45
 - validity: epilogue 49
- Manly-Parr estimates: survivorship (damselflies) 92–93
- Mann-Whitney U-tests
 - aggressive nest protection by waders 418–419, 426, 429
 - egg crypsis in blackbirds 167–168, 170
- Mannettia cordifolia* (Rubiaceae): *Aeria olenia* visits 106–107 Fig. 2 & Tab. 1
- mannose-6-phosphate isomerase (EC 5.3.1.8): *Bembicium vittatum* (snails) *Mpi* 62, 67 Tab. 2
- Mantel tests
 - RAPD variation in *Iliamna* 361
 - SE Asian shrews 197, 199, 202, 207–208, 215
- Marantaceae: pyrrolizidine alkaloids in ithomiine butterflies 106–107 Fig. 2 & Tab. 1, 110 Tab. 5
- Marantus* sp. 106–107 Fig. 2 & Tab. 1, 110 Tab. 5
- Marella*: Cambrian arthropods 23, 26
- martins
 - house (*Delichon urbica*): parasitic stress 474
 - purple (*Progne subis*): visual signals 391–392 Tab. 1
- *mass spectrometry: pyrrolizidine alkaloids (PAs) in ithomiine butterflies 99, 105
- Mata da Pedra (Sao Paulo, Brazil): pyrrolizidine alkaloids in ithomiine butterflies 101, 103, 108–109 Tab. 4 & Fig. 3, 112–114 & Tab. 9
- *mate choice: ornaments are armaments 385–386, 388–390
- material: first fossil anthomyid 442
- material and methods
 - complementary areas for diversity 128–132
 - evenness in moth populations 76–78 & Fig. 1
 - flower-fruit size allometry 402–404 & Fig. 1
 - pyrrolizidine alkaloids in ithomiine butterflies 101–105 & Fig. 1
 - RAPD variation in *Iliamna* 360–361
 - SE Asian shrews 199–203 & Fig. 1
 - specialization and speciation in shrimps 309–312
- *mating competition: ornaments are armaments 385–388
- *mating system evolution: monandry in damselflies 85–91 & Tab. 1, 95–96
- maxillopodans: arthropod evolution 13, 15–16
 - see also* *Bredocaris*
- mayflies: development of aquatic insect eggs *see* Ephemeroptera
- MDH-1 and MDH-2 (EC 1.1.1.37.)
 - Nucella lapillus* (dogwhelks) *Mdh-1* 180–181, 183 & Tab. 3, 185 Fig. 4
 - SE Asian shrews 200
 - MDH-1* 203
 - MDH-2* 204 Tab. 1, 206 Tab. 3
 - Mdh-2* 205 Tab. 2
 - sponge-dwelling alpheid shrimps
 - Mdh-1* and *Mdh-2* 310
 - Mdh-2* 313 Tab. 1
- Me*: SE Asian shrews 204–206 Tabs 1–3
- measurements
 - evenness in moth populations 76–78 & Fig. 1
 - growth similarities in tits 345
- **Mechanitis polymnia* (ithomiine butterflies): pyrrolizidine alkaloids 99–101, 104, 110 Tab. 5, 112 & Tab. 8, 115, 119
 - M. lysinnia* 110 Tab. 5
 - M. p. casabranca* 101
- Mecistoneuron*: fossil anthomyids 442
 - M. perpetuum* 442
- Mecistoneuron*: fossil anthomyids.
- Mediterranean
 - cladistic biogeography of the Eupreptantia (ME) 325, 327 Fig. 1, 329–335 & Tabs 1–2 & Figs 2–5, 338–340
 - see also* France, Italy and Sicily: *Brinkmannia*, *Drepanogigas*, *Paradrepanophorus* and *Pannettia*

- genetics of ice ages 249, 251–252
- Melungia salonina*: ithomiine visits 110 Tab. 5
- Melanochromis*: genetics of ice ages
- M. auratus* (cichlid mbuna) 275 App.
- M. heterochromis* 275 App.
- Melanotaeniidae* (rainbow fish): molecular data 276 App.
- Melospiza* (sparrows)
- genetics of ice ages 264
- M. melodia* (song sparrows): stress 473
- Merostomata: arthropod evolution 32, 41 Fig. 4
- Mesocapnia oenone* (Capniidae): development of aquatic insect eggs 224 Fig. 1, 238 App. 2
- Mesozoic: arthropod evolution 33–34
- *metabolic adaptation: development of aquatic insect eggs 221–223
- *metabolic potential: conservation strategies, stress and diversity 471, 473–477
- Metazoa: arthropod evolution 44
- methods
- aggressive nest protection by waders 419–418 & Figs 1–2
- cladistic biogeography of the Eurentantia 334–338
- development of aquatic insect eggs 226–228 & Fig. 3
- effect of light and diet on a polymorphic caterpillar 279–281
- egg crypsis in blackbirds 160–164 & Fig. 1
- geographic cohesiveness in an intertidal snail 59–63 Fig. 1
- growth similarities in tits 345–346
- molecular data (genetics of ice ages) 272–276 App.
- population genetic effects of a butterfly-ant mutualism 290–294 & Figs 1–2 & Tab. 1
- see also material and methods
- Methona themisto*: ithomiine visits 110 Tab. 5
- Mexico
- genetics of ice ages 249–250
- Quercus* (oaks) 178
- mice
- genetics of ice ages
- Mus*: hybrid zones 257
- see also *Mus*, *Onychomys* and *Peromyscus*
- stress
- oldfield *Peromyscus polionotus* 474
- white-footed (*Peromyscus leucopus noveboracensis*) 473
- microorganisms: mtDNA variability and male-lethal symbionts 453–456, 459, 462, 464, 467–468
- Mikania cordifolia* (Eupatoriae: Asteraceae): *Aeria olena* visits 106–107 Fig. 2 & Tab. 1, 110 Tab. 5
- Milankovitch theory: genetics of ice ages 248
- millipedes, pill (*Glomeris*): convergence 35
- *mimicry: *Nemoria arizonaria* (emerald moths) 277
- mimicry rings: convergence 35–36
- Miocene: closure of eastern Tethys Sea 340
- mitotypes: mtDNA variability and male-lethal symbionts 453–455, 468
- MOD (EC 1.1.1.40.): SE Asian shrews 200
- models
- conservation strategies, stress and diversity 471, 477
- development of aquatic insect eggs 236–237 App. 1
- effect of light and diet on a polymorphic caterpillar: I ANOVA 281
- evenness in moth populations 75, 77–78, 81–83
- see also Broken Stick (BS), Overlapping Niche (ON) and Sequential Break (SB)
- multivariate genetic differentiation 173
- multivariate 175–180 & Fig. 1 & Tab. 1, 187
- ornaments are armaments 385–387, 389–390, 393–394
- see also Fisher run-away and indicator models 386
- population genetic effects of a butterfly-ant mutualism 287–288
- quantitative: mtDNA variability and male-lethal symbionts 455–459, 463–464, 467
- SE Asian shrews 197, 202, 208–212 & Fig. 6, 214–216
- simulation: genetics of ice ages 250
- statistics of synapomorphies 484
- Molara*: Cambrian arthropod 26
- *molecular approaches: arthropod evolution 1–2, 13, 18, 25–26 & Fig. 1, 28–29
- limitations 44–47, 55
- molly, sailfin (*Poecilia velifera*): visual signals 391–392 Tab. 1
- Molothrus ater* (brown-headed cowbirds): ornaments are armaments 390, 391–392 Tab. 1
- monandry: damselfly evolution 85–89, 93–94, 96–97
- Monjolinho Arboretum (San Paulo, Brazil):
- pyrrolizidine alkaloids in ithomiine butterflies 101, 108–110 & Tabs 3 & 5
- *monophyly: arthropod evolution 1, 34–37
- monophyly polyphyly debate 34–37 & Fig. 3, 44
- Monotremata: statistics of synapomorphies 490–492 & Tabs 2–3, 496–498 Apps. 1–2 & Tab. A4
- Montastrea annularis*: coral reefs 321
- Monura: arthropod evolution 8
- Moran's I: multivariate genetic differentiation 192
- *morphology: growth similarities in tits 343, 350 Fig. 1, 352
- see also measurements, size and traits
- mosquitoes see Culicidae
- moths
- Arctiidae: pyrrolizidine alkaloids 99–100, 114, 117–118
- cornblower (*Otrinia nubilalis*: introduction to USA 263
- effect of light and diet on a polymorphic caterpillar 277–283 & Fig. 2
- hawkmoths (Sphingidae) 278
- phycitid 278
- evenness and species numbers 75–84 & Figs 1–3 & Tab. 1
- Greya*
- molecular data 274 App.
- switch of umbrellifer hosts 266
- Heliothis*: mtDNA variability and male-lethal symbionts 467
- whistling (*Hecatesia thyridion*): acoustic signals 391–392 Tab. 1
- MPI (EC 5.3.1.8.)
- SE Asian shrews 200, 203
- Mpi* 204–206 Tabs 1–3
- sponge-dwelling alpheid shrimps: *Mpi* 310, 312–313 Fig. 1 & Tab. 1
- mtDNA: stress, metabolic potential and conservation 476
- *mtDNA variability: male-lethal symbionts 453–470 & Figs 1–6 & Tabs 1–3 & App
- Mulinia lateralis* (coot clam): heterozygosity 474

- multiple correspondence analysis (MCA): multivariate genetic differentiation 174–179
see also constant row total-multiple correspondence analysis (CRT-MCA)
- *multivariate analyses: genetic differentiation 173–178, 194–195 App. & Fig. 8, 182–187 & Tab. 3 & Figs 3–6
 multivariate analysis of variance (MANOVA) 175, 191–192 & Tab. 4
- Muricea (*Nucella lapillus*; dogwhelks): multivariate genetic differentiation 175
- Mus (mice): genetics of ice ages 257, 259
M. castaneus 275 App.
M. domesticus 275 App.
M. musculus (housemice) 275 App.
- Muscidae: Diptera 443, 446–447
- mussels
 coral-dwelling 321
Mytilus edulis: heterozygosity 474
- mutations: mtDNA variability and male-lethal symbionts 454–457, 460, 462–466 & Tab. 1, 468
- Mutisidae, Asteraceae: visits by Ithomiinae 109
see also *Gochmatia*
- *mutualism, butterfly-ant: population genetic effects 287–290
- Myodocopina: compound eyes 41–42 & Fig. 4
- Myriapoda: arthropod evolution 2, 5, 19, 34 Fig. 3, 41 Fig. 4
- *myrmecophily: population genetic effects of a butterfly-ant mutualism 287–289, 300, 303
- Myscelia orsis* (nymphalines): experimental spider prey 105
- Mysidopsis* (malacostracan): arthropod evolution 7
- Mystacocarida: eyeless or without compound eyes 41–42 & Fig. 4
- Mytilus*
M. edulis (mussels)
 heterozygosity 474
 localised selection 58
 mtDNA variability and male-lethal symbionts 467
- Nally, Ralph C. Mac. *Ecological versatility and Community Ecology*. (Book Review: Mark O. Hill) 245
- Naraia*: eyelessness 42
- *narrow endemism: complementary areas for diversity 126–127, 130, 132, 138, 144–146 App. 1
- Matrix* (grass snakes): hybrid zones 257
- Naupeta cinerea* (cockroaches): ornaments are armaments 388
- nauplii: arthropod evolution 3–4, 12–13, 16, 19–23
- Nebalia*: malacostracan crustacean 16
- Nei, M.
 diversity analysis: geographic cohesiveness in an intertidal snail 68
 genetic distances (D)
 population genetic effects of a butterfly-ant mutualism 287, 293–294, 296–298 & Tab. 4 & Figs 3–4, 300 Fig. 5
 SE Asian shrews 201, 203, 208–209 Tab. 4, 213
 specialization and speciation in shrimps 310, 312 Fig. 1, 314
- Nematocera: *Nephila* diet 120
- *Nemertea: cladistic biogeography of the Eurentantia 325–332 & Fig. 1, 335–336 Figs 4–5, 338, 340
- **Nemoria arizonaria* (emerald moths): effect of light and diet on a polymorphic caterpillar 277–283 & Fig. 2
- Nemoria* 283
- Nemouridae (Plecoptera): development of aquatic insect eggs 230, 238–239 App. 2
- Nemoura arctica* 238 App. 2
N. avicularis 238 App. 2
N. cambria 238 App. 2
N. cinerea 228, 238 App. 2
N. erratica 239 App. 2
N. pictetti 228
N. viki 239 App. 2
Nemurella pictetti 239 App. 2
see also *Amphinemura*, *Malenka* and *Zopada*
- neoplatyphylline: pyrrolizidine alkaloid 114
- Neopolina*: metamerism 44
- Neotoma* (woodrats): genetics of ice ages 261, 264
N. albigula: molecular data 274 App.
N. floridana (eastern woodrats) 261, 274 App.
N. micropus 274 App.
- Neotropics
 complementary areas for plant diversity 125, 128, 131
see also Caryocaraceae, Chrysobalanaceae, Dichapetalaceae, Lecythidaceae and Proteaceae
- first fossil anthomyid clade 441, 446
- **Nephila clavipes* (orb weaving spiders): predation of ithomiine butterflies 99–101, 105, 115
 chemical defences against *Nephila* 118–121
- Nerita*: intertidal communities
N. atramentosa 61, 64 Tab. 1
N. plicata 61, 64 Tab. 1
- Nerophis ophidion* (pipefish): visual signals 391–392 Tab. 1, 394
- Neruda* (butterflies): molecular data 274 App.
- nesting habits: growth similarities in tits 344–345, 347, 349
 hole-nesting 344–345, 353
 open-nesting (finches) 344–345, 353
- nests
 egg crypsis in blackbirds 159–163 & Fig. 1, 167–170 & Fig. 4
 habitats: aggressive protection by waders 409–411, 414–416, 419, 423–424 & Tab. 1, 426, 429, 436–439 App.
- *neutrality: geographic cohesiveness in an intertidal snail 57
- New Caledonia, Western Pacific (WP): Eurentantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
- New Guinea, Western Pacific (WP): Eurentantia 327–328 & Fig. 1, 335 Fig. 4, 339
- New Mexico: *Vulpes* molecular data 274 App.
- New South Wales, Australia: butterfly-ant mutualism 287–288, 290–292 & Fig. 1 & Tab. 1
- New Zealand and Australia: monochromatic *Ischnura aurora* females 96
- newts (*Triturus*): genetics of ice ages 264
 crested (*T. cristatus*) 260–261
- Nodilittorina*: geographic cohesiveness in intertidal snails
N. australis 61, 64 Tab. 1
N. pyramidalis 61, 64 Tab. 1
N. unifasciata 61, 64 Tab. 1
- North America
Crepis tectorum 402

- genetics of ice ages 247–252, 257, 260–262, 272–273 App.
 monandry in damselflies 86
 refugia 198
 Western Atlantic Ocean (WAT): Eureptantia 327
 Fig. 1, 329
see also Canada and U.S.A.
- Norway, Eastern Atlantic (EAT): Eureptantia 329
- Notodromas*: development of ocellus 42
- Notonemouridae (Plecoptera): development of aquatic insect eggs 230, 239 App. 2
see also *Austrocerella*
- Notostraca: arthropod evolution 3, 13–14
- Nousia* (Leptophlebiidae): development of aquatic insect eggs
N. fuscula 241 App. 2
N. inconspicua 241 App. 2
- **Nucella lapillus* (dogwhelks): multivariate genetic differentiation 173, 175, 180–182 & Fig. 2, 187–190
- Nucella lapillus*: localised selection 58
- null hypotheses
 flower-fruit size allometry 401, 403–405
 RAPD variation in *Iliamna* 360, 364
- numbers (parents): aggressive nest protection by waders 409, 411–412, 414, 418, 420 Fig. 4, 423–429 & Tabs 1–2, 436–439 App.
- Numenius*: aggressive nest defence
N. americanus 437 App.
N. arquata 437 App.
N. borealis 413, 437 App.
N. madagascariensis 413, 437 App.
N. minutus 437 App.
N. phaeopus (whimbrels: Hudsonian curlews) 411, 437 App.
N. tahitiensis 437 App.
N. tenuirostris 413, 437 App.
- Nymphalidae: polymorphic caterpillars 278
- Nymphalinae: pyrrolizidine alkaloids in ithomiine butterflies 109, 119
- Nymphalidae 99, 109
see also *Biblis* and *Myseelia*
- oaks *see* *Quercus*
- Obi (Indonesian archipelago): Eureptantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
- Oceanapidae (sponges) 321
Oceanapia sp. 316, 318 Fig. 4, 321
- Odocoileus virginianus* (white-tailed deer): molecular data 274 App.
- *Odonata: development of aquatic insect eggs 221, 226–233 & Figs 3–5, 240 App. 2
see also Coenagrionidae, Corduliidae and Libellulidae
- *Odonata: monandry in damselflies 85, 90, 96
- Oeciacus hirundinis* (house martin bugs): ectoparasite 474
- Oland, Baltic island: *Crepis tectorum* subsp. *pumila* 402
- Oligocene: *Mecistoneuron perpetuum* 442, 451
- ommatidia: compound eyes 37–39
- Onchorhynchus mykiss* (rainbow trout): bacterial gill disease 474
- *ontogeny: growth similarities in tits 343–346, 349–354 & Tabs 3–5 & Fig. 1
- Onychomys* (grasshopper mice): genetics of ice ages 261
- O. aremicolor*: molecular data 273 App.
O. leucogaster 273 App.
O. torridus 273 App.
- Onychophora: arthropod evolution 2, 5, 11, 15, 41–43 & Fig. 4
 monophyly 45
- Opabinia*: arthropod evolution 12, 37
- Ophrysopus* sp. (Asteraceae): pyrrolizidine alkaloids in ithomiine butterflies 108 Tab. 3, 110 Tab. 5, 113 & Tab. 9, 117
- Opsanus* (toadfish): genetics of ice ages 264
O. beta: molecular data 272 App.
- orders: development of aquatic insect eggs 221, 228–229, 233
- Ordovician: arthropod evolution 29–30, 42
- ornaments: an evolutionary explanation 385–394 & Tab. 1
- Orsten fossils: arthropod evolution 23, 26
- Orstencarida: compound eyes 41 Fig. 4
- Oryctolagus cuniculus* (European rabbits): genetics of ice ages 260
- Osmerus* (smelt): genetics of ice ages 261
O. dentex: molecular data 273 App.
O. eperlanus 273 App.
O. mordax 273 App.
- Ostracoda: eyeless or without compound eyes 41–42 & Fig. 4, 190
- Ostrinia nubilalis* (cornblower moths): introduction to USA 263
- outgroups: cladistic biogeography of the Eureptantia 330–332, 334–338 & Figs 3–5
- overlapping niche (ON): evenness in moth populations 75, 77–78 & Fig. 1, 82
- Ovis*
O. canadensis
 bighorn sheep: ram horn volume 475
 mountain sheep: visual signals 391–392 Tab. 1
O. dalli (Nearctic wild sheep): genetics of ice ages 260
- Oxalidaceae: *Aeria olena* visits 106–107 Fig. 2 & Tab. 1
Oxalis rhombocaula 106–107 Fig. 2 & Tab. 1
- oystercatchers: aggressive nest defence 422, 427
- oysters, American (*Crassostrea virginica*): molecular data 272 App.
- PA (EC 3.1.3.2.): SE Asian shrews 200, 204–206 Tabs 1–3
Pa 206 Tab. 3
- Pacific
 cladistic biogeography of the Eureptantia 325, 327
 Eastern (EP) 330, 333, 340
 Western (WP) 327–329 & Fig. 1, 331–337 & Tab. 1 & Figs 2–6, 339–340
see also Australia (Gbr), New Caledonia and New Guinea: *Drepanophoresta*, *Drepanophorina*, *Pummettia*, *Urichonemertes* and *Xenonemertes*
 genetics of ice ages 249, 272 App.
- Padogobius martensi* (fish): ornaments are armaments 389–390
- Palaeartic regions: SE Asian shrews 197, 203, 213
- Palaeocene: eureptantian phylogenetic trees 339
- palaeoclimates: genetics of ice ages 247–248, 262
- Palaeozoic: arthropod evolution 8–9, 14, 21, 33–34, 49
- Palawan: SE Asian shrews 200 Fig. 1, 209, 214

- Palicourea* sp. (red flowers: Rubiaceae): *Tithorea harmonia* visits 108–109 & Tab. 4 & Fig. 3
- Panama
- Nephila clavipes* webs 120
- sponge-dwelling alpheid shrimps 307–321 & Figs 1–5 & Tabs 1–2
- Panopsis*: complementary areas for diversity 128–129, 132–133 Fig. 1
- P. cinnamomea* 147 App. 2
- P. ferruginea* 151 App. 2
- P. macrocarpa* 147 App. 2
- P. ptariana* 147 App. 2
- P. rectistylus* 150 App. 2
- P. yolombo* 148 App. 2
- P. yungasensis* 151 App. 2
- Papilionidae (swallowtail butterflies): colour polymorphism 278
- Papilioninae
- Papilionidae: eaten by *Nephila* 119
- Parides* spp.: non-pyrrolizidine alkaloid insects 109
- Paradrepanophorus*: eurentatic nemerteans 339
- P. corallinicola* (ME) 327 Fig. 1, 329, 332 Fig. 2
- P. crassus* (ME, EAT) 327 Fig. 1, 329, 332 Fig. 2, 339
- P. nissidensis* (ME) 327 Fig. 1, 329, 332 Fig. 2
- P. obiensis* (EIO: Obi) 327–328 & Fig. 1, 332 Fig. 2
- *parental care systems: aggressive nest protection by waders 409–411, 414, 418, 426–427, 436–439 App.
- Parides* spp. (Papilioninae): non-pyrrolizidine alkaloid insects 109
- *parsimony: cladistic biogeography of the Eurentantia 325–326, 328–338 & Figs 2 & 5–6 & Tab. 1, 338, 340
- see also BPA
- parsimony principle: statistics of synapomorphies 483–484
- parsonsine: isolation from Echinoidea (Parsonsia) 110
- partridges, grey (*Perdix perdix*): visual signals 391–392 Tab. 1
- Parus*
- genetics of ice ages: North American chickadees 261, 264
- P. atricapillus* 273 App.
- P. carolinensis* 273 App.
- growth similarities in tits
- P. coerules* (blue tits) 343
- P. major* (great tits) 343
- ornaments are armaments: *P. major* (great tits) 387, 391–392 Tab. 1
- Passer domesticus* (house sparrows): ornaments are armaments 387, 391–392 Tab. 1
- Passerella* (fox sparrows): growth
- P. i. iliaca* 352
- P. i. megarhyncha* 352
- passerines: egg crypsis in blackbirds 159
- patterns
- alkaloids (PAs) in ithomiine butterflies 99, 110, 112, 114, 117–118
- cladistics of the Eurentantia 325–326, 330, 334
- complementary areas for diversity 126
- development of aquatic insect eggs 225, 232
- egg crypsis in blackbirds 159–161 & Fig. 1
- brightness disparity (BD) 159, 164–170 & Tabs 1–3 & Figs 2–4
- mean brightness (MB) 159, 164–168 & Tabs 1–3 & Figs 2–4
- pattern disparity (PD) 159, 164–168 & Tabs 1–3 & Figs 2–4, 170
- geographic cohesiveness in an intertidal snail 57–59, 62, 64–65, 67–72
- growth similarities in tits 343–345, 347–350, 352–354
- host use: sponge-dwelling alpheid shrimps 309, 320–322
- monandry in damselflies 86–88, 91 Tab. 1, 93–97
- multivariate genetic differentiation 173–175, 181, 183–185 & Fig. 5, 191 Tab. 4
- population genetic effects of a butterfly-ant mutualism 287–288, 301
- SE Asian shrews 197, 199, 203, 214, 216
- statistics of synapomorphies 483, 486–491 & Tabs 1–3, 495–496 & Tabs A1–A3, 495–496 App. 1: Tabs A1–A3
- PAUP programs
- cladistic biogeography of the Eurentantia 328–329
- SE Asian shrews 201
- Pauropoda: eyeless or without compound eyes 41 Fig. 4
- Pavo cristatus* (peacocks): visual signals 391–392 Tab. 1
- Pax-6* (control genes): eye development 54
- peacocks (*Pavo cristatus*): visual signals 391–392 Tab. 1
- Pelsaert Group (Houtman Albrolous Islands, Western Australia): *Bembicium vittatum* (snails) 60 Fig. 1, 63, 65 & Fig. 2, 67–72 & Tab. 2 & Fig. 4
- penes
- monandry in damselflies 85–86, 88, 91 Tab. 1, 95, 97
- Symphetrum danae* (anisopteran) 95
- Peninsula Malaysia (PM): SE Asian shrews 197–198, 200 Fig. 1, 202–204 & Fig. 2 & Tab. 1, 208–212 & Figs 5–6, 215, 219 App.
- 'Continent' 210, 212 Fig. 6
- Pep-1* & *Pep-2*, allozyme variations (*Nucella lapillus*: dogwhelks): multivariate genetic differentiation 180–181, 183 Tab. 3, 185 Fig. 4
- Pep-2* 183
- Perdix perdix* (grey partridges): visual signals 391–392 Tab. 1
- Perlidae (Systellognatha): development of aquatic insect eggs 239 App. 2
- Perla bipunctata* 239 App. 2
- P. burmesiana* 230
- see also *Dinocras*
- Perlidae (Systellognatha): development of aquatic insect eggs 239 App. 2
- Perlodes microcephala* 230, 239 App. 2
- see also *Diura*, *Isogenus*, *Isoperla* and *Skwala*
- Permian
- arthropod evolution 1–2, 5, 9, 33
- Ice Age: Plecoptera 229
- Peromyscus*, *P. maniculatus* (deer mice): molecular data 274 App.
- stress
- P. leucopus noveboracensis* (white-footed mice) 473
- P. polionotus* (oldfield mice) 474
- Peters Mountain (PM), Virginia: *Ilamma* 358–365 & Tabs 1–3
- Petrogale* (rock wallabies): molecular data

- P. assimilis* 276 App.
P. herberti 276 App.
P. purpureicollis 276 App.
Petrosiidae (sponges) 321
 see also *Xestospongia*
Pgd see phosphogluconate dehydrogenase (EC 1.1.1.44): *Bembicium vittatum* (snails)
6PGD (EC 1.1.1.42.): SE Asian shrews 200, 204–206
 Tabs 1–3
PGI (EC 5.3.1.9.)
 SE Asian shrews 200, 204–205 Tabs 1–2
 sponge-dwelling alpheid shrimps: *Pgi* 310, 313–314
 & Tab. 1
 Pgm: sponge-dwelling shrimps 310, 313 Tab. 1
 PGM (EC 2.7.5.1.)
 SE Asian shrews 200
 Pgm 204–206 Tabs 1–3
 see also phosphoglucomutase
Pgm-1 (EC. 5.4.2.2.): *Jalmenus evagoras* 291, 295 Tabs
 2–3
 Pgm-1 299 Tab. 5
Pgm-1 and *Pgm-2*: *Nucella lapillus* (dogwhelks) 180–181,
 183 Tab. 3, 185 Fig. 4
Pgm-1 184
Phalaropus: aggressive nest defence 427
 P. fulvicaeus 438 App.
 P. lobatus 438 App.
 P. tricolor 438 App.
Phaenantho: first fossil anthomyid and comments on
 clade definitions 441, 445–448 & Fig. 3, 450–451
 & Fig. 5
Phasianus colchicus (pheasants): visual signals 391 392
 Tab. 1
pheasants (*Phasianus colchicus*): visual signals 391 392
 Tab. 1
Phaidole spp. (ants): seed removal 303
Pherbellia cinerella (Sciomyzidae): development of
 aquatic insect eggs 242 App. 2
pheromone precursors: alkaloids (PAs) in ithomiine
 butterflies 99–100
Philippines
 SE Asian shrews 197, 200 Fig. 1, 202–204 & Fig. 2
 & Tab. 1, 207–212 & Figs 5–6, 214–216, 219
 App.
 subarea (Eastern Indian Ocean, EIO): Eureptantia
 328
Philomachus pugnax: aggressive nest defence 439 App.
Phospatocopida
 ?compound eyes 41 Fig. 4
 labrum 16
phosphoglucomutase (EC 2.7.5.1)
 Bembicium vittatum (snails)
 Pgm1 65, 72
 Pgm1 and *Pgm2* 62, 67 Tab. 2
 see also PGM
phosphogluconate dehydrogenase (EC 1.1.1.44):
 Bembicium vittatum (snails) *Pgd* 62, 65, 67 Tab. 2, 72
phosphoglucose isomerase (PGI): stress 476
 pythid moths: colour polymorphism 278
Phyllocarida: compound eyes 41 Fig. 4
Phylodiaptomus annae: arthropod evolution 4
*phylogeny: SE Asian shrews 197, 201–203 & Fig. 2,
 206–209, 212–216 & Fig. 5
phylogeny
 Eureptantia 326–327 & Fig. 1, 330
 first fossil anthomyid and comments on clade
 definitions 441, 447–451 & Figs 4–5
 *phylogeography: genetics of ice ages 247, 257, 262
 *phylogeography: mtDNA variability and male-lethal
 symbionts 453
Phymosia remota: *Iliamna corei* (Peters Mountain mallow)
 358–359
Physiphora demandata (flies): ornaments are armaments
 389
Pieridae: polymorphic caterpillars 278
 Pieris brassica 278
Pierini: pyrrolizidine alkaloids 120
Pieridae 119
pigmentation quantification: egg crypsis in blackbirds
 168–169
pines
 lodge pole: genetics of ice ages 260
 pollen cores: Europe 250
 spruce budworm: mtDNA divergence 266
Pinus taeda (loblolly pines): moisture stress 479
 pipefish (*Nephris ophidion*): visual signals 391–392 Tab.
 1, 394
Placidula euryanassa (ithomiine): *Brugmansia suaveolens*
 (host plant) 118
plankton, freshwater: copper as a stress 479
*plant taxonomy: RAPD variation in *Iliamna* 357
plants
 genetics of ice ages 250–252
 see also alder, beech, grass, hazel and pines
 host: population genetic effects of a butterfly-ant
 mutualism 287–290, 292 & Tab. 1, 300–303
 see also *Acacia*
 inbreeding depression 473, 476
Plasmodium mexicanum: malarial parasite (lizards) 174
Platypezidae: *Mecistoneuron perpetuum* 442
 platyphylline: pyrrolizidine alkaloid 114
*Plecoptera (stoneflies): development of aquatic insect
 eggs 221, 226–230 & Figs 3–5, 232–233,
 238–240 App. 2
 see also Eutholognatha and Systellognatha
Pleistocene
 genetics of ice ages 247–249, 257–258, 261–262,
 264–265 & Fig. 7
 SE Asian shrews 198–200 & Fig. 1, 209, 211, 213,
 216
*plesion: first fossil anthomyid and comments on clade
 definitions 441, 449–450
Plethodon cinereus (salamanders): genetics of ice ages 260
Pliocene: genetics of ice ages 248, 262
plovers (*Pluvialis*): aggressive nest defence 427
 American golden (*P. dominica*) 426, 436 App.
 crab 427
 Eurasian golden (*P. apricaria*) 427–428, 436 App.
 golden spp. 426
 grey (*P. squatarola*) 411, 422, 436 App.
Pluvialis (plovers): aggressive nest protection by waders
 P. apricaria (Eurasian golden) 427–428, 436 App.
 P. dominica (American golden) 426, 436 App.
 P. fulva 436 App.
 P. squatarola (grey) 411, 422, 436 App.
Poaceae (*Buchloe dactyloides*): use of AMOVA 360
Podocopida: development of ocellus 42
Poecilia: ornaments are armaments
 P. latipinna 389
 P. reticulata 391–392 Tab. 1

- P. velifera* (sailfin molly) 391–392 Tab. 1
- Poecilia reticulata* (guppies): molecular data 272 App.
- poikilotherms: development of aquatic insect eggs 223, 232
- pollen records: genetics of ice ages 249–250, 252, 262
- polyandry: damselfly evolution 85–89, 94, 96
- polychromatism: damselflies 91, 96
- *polyphenism: *Nemoria arizonaria* (emerald moths) 277
- *polyphyly: arthropod evolution 1, 25–34 & Figs 1–2
- monophyly polyphyly debate 34–37 & Fig. 3, 47
- Popperian argument: statistics of synapomorphies 484, 493
- *population structure: genetics of ice ages 247–248, 251–255 & Figs 1–2, 257–261, 264, 266, 272–274 App.
- populations
- Bembicium vittatum* (snails) 57–59, 68, 71–73
 - conservation strategies, stress and diversity 471–473, 475 477, 479
 - Crepis* (Asteraceae) 401–406
 - development of aquatic insect eggs 221, 223, 226–228 & Fig. 3, 230 Fig. 5, 232
 - genetics: effects of a butterfly-ant mutualism 287–303 & Figs 1–4 & Tabs 2–5
 - mtDNA variability and male-lethal symbionts 453–456, 458–68 & Figs 4–6 & Tabs 1–3
 - multivariate genetic differentiation 173–195 & Figs 1–5 & Tabs 1–4 & App.
 - RAPD variation in *Ikhanna* 357–361 & Fig. 1, 364–365 & Fig. 2 Tab. 3
 - James River 358 & Fig. 1, 360
 - Mallow Preserve (MP) 358 Fig. 1, 360, 364–365 & Fig. 2 & Tab. 3
 - see also Interstate 64(I), Iron Gate (IG) and Peters Mountain (PM)
 - SE Asian shrews 197, 202 203, 206, 208–209 & Tab. 4, 215
 - sponge-dwelling alpheid shrimps 307–315 & Fig. 1 & Tab. 2, 319–321
- pratincoles: aggressive nest defence 427
- see also *Glareola*
- Prawle Point (Devon coast): *Nucella lapillus* (dogwhelks) 181–187 & Figs 2–3 & 5–6, 189–190
- prawns, decapod: Atyidae: arthropod evolution 4
- Pre-crustaceans: eyeless or without compound eyes 41 Fig. 4
- Precambrium: eurentatic phylogenetic trees 339
- Precis coenia* (Lepidoptera): arthropod evolution 7
- predation
- egg crypsis in blackbirds 159–162 & Fig. 1, 167–168 & Fig. 4
 - orientation
 - chemical: spiders (*Nephila clavipes*) 99–101, 105, 120–121
 - visual: birds 120
 - regimes: aggressive nest protection by waders 409–414, 416–417, 420, 422–424 & Tab. 1, 426–429, 436–439 App.
- presentation, nonhistoric: monandry in damselflies 86 87
- Prestonia* (Apocynaceae: Echitoideae): ithomiine foodplants
- P. acutifolia* 99, 101, 108–111 & Tabs 4 & 6 & Fig. 3, 118
 - P. coalita* 101, 111, 118–119
- prey, alternative: aggressive nest protection by waders 409, 411–412, 417, 423–426 & Tabs 1–2, 428–429, 436–439 App.
- principal components analyses
- geographic cohesiveness in an intertidal snail 61–65 & Tab. 1 & Fig. 2, 67
 - monandry in damselflies 88–89
 - multivariate genetic differentiation 173, 179, 191 Tab. 4
- principal coordinates analysis (PCoA): multivariate genetic differentiation 174–175, 191
- Prituitzia hymenaea*: ithomiine visits 110 Tab. 5
- probabilities
- mtDNA variability and male-lethal symbionts 455–459
 - statistics of synapomorphies 485–489 & Tab. 1 & Fig. 2, 492–493
 - see also Freeman-Tukey method
- Proboscidae (elephants): convergence 36
- Procavia capensis* (rock hyrax): molecular data 275 App.
- Progne subis* (purple martin): visual signals 391–392 Tab. 1
- pronghorns: ornaments are armaments 389
- Prosobranchia (*Nucella lapillus*, dogwhelks): multivariate genetic differentiation 175
- PROT and XDH (EC 1.2.3.2.): SE Asian shrews 200
- PROT 203
- see also XDH
- Proteaceae: complementary areas for diversity 128–129, 132–138 & Figs 1 & 3–4 & 6–7
- see also *Panopsis*
- protomandibulates: arthropod evolution 19–20
- Protonemura* (Nemouridae): development of aquatic insect eggs
- P. meyeri* 239 App. 2
 - P. montana* 239 App. 2
 - P. praecox* 239 App. 2
- Protura: eyeless or without compound eyes 41 Fig. 4, 43
- pruinescence: monandry in damselflies 85, 87, 91 Tab. 1, 94–96
- Pseudocada erruca*: ithomiine visits 110 Tab. 5
- Pseudotrophus zebra*: molecular data 275 App.
- Psychotria* (Rubiaceae): pyrrolizidine alkaloids in ithomiine butterflies
- P. leiocarpa* 110 Tab. 5
 - P. melaneoides* 108 Tab. 3
- ptarmigans, rock (*Lagopus mutus*): visual signals 391–392 Tab. 1
- Pterygota: compound eyes 41 Fig. 4, 43
- Pummetia*: eurentatic nemerteans
- P. terina* (EIO: Ambon) 327–328 & Fig. 1, 332 Fig. 2
 - P. hubrechtii* (ME) 327 Fig. 1, 329, 332 Fig. 2
 - P. maldivensis* (EIO: Timor, WIO: Maldives) 327 328 & Fig. 1, 332 Fig. 2, 339
 - P. micrommata* (WP: New Guinea) 327–329 & Fig. 1, 332 Fig. 2, 339
 - P. spectabilis* (ME, EAT) 327 Fig. 1, 329, 332 Fig. 2
 - P. splendida* (ME, EAT) 327 Fig. 1, 329, 332 Fig. 2, 339
 - P. timorensis* (EIO: Timor) 327 328 & Fig. 1, 332 Fig. 2
 - P. willeyana* (EIO: Seram, WIO: Maldives) 327–328 & Fig. 1, 332 Fig. 2, 338–339

- pupae
 butterfly-ant mutualism 289–290
 effect of light and diet on a polymorphic caterpillar 278, 283
- pupfish, Pecos (*Cyprinodon pecosensis*): visual signals 391–392 Tab. 1
- Pyrenees: genetics of ice ages 249, 251–253, 258–259 & Fig. 5, 262–266 & Fig. 7
- pyrrolizidine alkaloids (PAs): ithomiine butterflies 99–123 & Figs 1–3 & Tabs 1–11
see also anobiline, echinatine, heterophylline, indecine, intermedine, lycopsamine, parsonsine, rinderine and supinine
- Quercus* (oaks): effect of light and diet on a polymorphic caterpillar
 catkins 277–283 & Tab. 1 & Fig. 2
 leaves 277, 279–283 & Tab. 1 & Fig. 2
Q. arizonica 278
Q. emoryi (emory oaks) 278–280
 twigs 277, 279
- 'R' program: SE Asian shrews 202
- rabbits, European (*Oryctolagus cuniculus*): genetics of ice ages 260
- *radiation: arthropod evolution 1, 22–24, 26, 47
- radiation
 SE Asian shrews 197, 211
 sponge-dwelling alpheid shrimps 307–309, 321
- *random amplified polymorphic DNA: variation in *Ilamna* *see* RAPD
- *range changes: genetics of ice ages 247–248, 251–264 & Fig. 1, 266
- *RAPD (random amplified polymorphic DNA): variation in *Ilamna* 357, 359–365 & Tabs 1–2 & Fig. 3
- Rapoport's Rule: species diversity 263
- *rarity: complementary areas for diversity 126, 130–134 & Fig. 4 & Tab. 1, 137 Tab. 2, 144–148 App. 1, 155–157 App. 4
 rarity scores 131–132, 134–135, 155–157 App. 4
- rats, mole (*Spalax ehrenbergi*)
 aridity stress 474–476
 molecular data 275 App.
- Recent: arthropod evolution 23, 30
- *recolonization: population genetic effects of a butterfly-ant mutualism 287, 289, 300–303
- Recurvirostra* (avocets): aggressive nest defence
R. americana 436 App.
R. avosetta 436 App.
- redshank (*Tringa totanus*): aggressive nest defence 426, 437 App.
- refugia: Amazon basin and North America 198, 215
- *refugia: genetics of ice ages 247, 250–256 & Fig. 3, 258–265 & Figs 5–7
- regressions
 development of aquatic insect eggs 224, 227–229, 236–237 App. 1
 evenness in moth populations 75, 82
 geographic cohesiveness in an intertidal snail 67–70 & Tab. 3
 growth similarities in tits 346–347 & Tab. 4
 major axis: flower-fruit size allometry 401–402, 404
 stepwise: aggressive nest protection by waders 422–427 & Tabs 1–2
 wing lengths in damselflies 90 Fig. 1
- Rehbachella*: Cambrian arthropod 3–4, 16, 54–55
- Remipedia*: crustacean 4, 41 & Fig. 4
- reptiles: ornaments are armaments 387
see also lizards
- *resource apportionment: evenness in moth populations 75
- results
 aggressive nest protection by waders 418–425 & Figs 3–6 & Tabs 1–2
 butterfly-ant mutualism 294–303 & Tabs 2–5 & Figs 3–5
 cladistic biogeography of the Eurentantia 331–333 & Fig. 1 & Tabs 1–2
 complementary areas for diversity 134–136 & Figs 5–7 & Tab. 1–2
 development of aquatic insect eggs 228–229
 effect of light and diet on a polymorphic caterpillar 281–282 & Tab. 1
 egg crypsis in blackbirds 164–168 & Tabs 1–3 & Figs 2–4
 evenness in moth populations 78–81 & Figs 2–3 & Tab. 1
 flower-fruit size allometry 404–405 & Fig. 2
 genetic analysis of shrimp populations 312–318 & Figs 1–4 & Tabs 1–2, 320
 geographic cohesiveness in an intertidal snail 63–70 & Tabs 1–4 & Figs 3–5
 growth similarities in tits 346–349 & Tabs 1–2
 mtDNA variability and male-lethal symbionts 459–464 & Figs 1–4
 pyrrolizidine alkaloids in ithomiine butterflies 105–115 & Figs 2–3 & Tabs 1–9
 RAPD variation in *Ilamna* 361–365 & Tabs 1–3 & Figs 2–3
 SE Asian shrews 203–213 & Tabs 1–4 & Figs 3–6
 retention indices (chromatography): pyrrolizidine alkaloids in ithomiine butterflies 105
- RFLPs (restriction fragment length polymorphisms): genetics of ice ages 257, 261, 272–276 App.
- Rhagoletis pomonella* (hawflies): genetics of ice ages 266
- Rhithrogena* (Heptageniidae): development of aquatic insect eggs
R. cf. hybrida 241 App. 2
R. loyolae 221, 228–233 & Fig. 5, 241 App. 2
R. semicolata 228, 231, 241 App. 2
- Rhizocephala, parasitic: arthropod evolution 41 Fig. 4
- Rhytidoponera* spp. (ants): seed removal 303
- Ricinulei: blindness 42
- rinderine (pyrrolizidine alkaloid): ithomiine butterflies 99, 102 Fig. 1, 104, 109–110, 113–114, 117
- RNA sequences: arthropod evolution 44–46
- rodents
 molecular data: 'murid' (*Akodontine* tribe) 274 App.
 population fluctuations: aggressive nest protection by waders 412, 417, 423, 428–429
- Roger's distances: SE Asian shrews 201–202, 208–210 Tab. 4 & Fig. 3
- Rosaceae: *Aeria olena* visits 106–107 Fig. 2 & Tab. 1
Eribotrya sp. 106–107 Fig. 2 & Tab. 1
- rosefinches, scarlet: open-nesting 344
- Rostratula benghensis*: aggressive nest defence 436 App.
- Rubiaceae: pyrrolizidine alkaloids in ithomiine butterflies 106–110 Figs 2–3 & Tabs 1 & 3–5

- see also* *Borreria*, *Coffea*, *Manettia*, *Palicourea* and *Psychotria*
- Ruellia longifolia* (Acanthaceae): *Aeria olena* visits 106–107 Fig. 2 & Tab. 1
- Russia: genetics of ice ages 249, 251, 258–259 & Fig. 5, 273 App.
- Rutaceae: pyrrolizidine alkaloids 110 Tab. 5
- Esenbeckia leiocarpa* 110 Tab. 5
- Ryan's Q test: morphometric analysis of shrimp populations 315 Tab. 2
- salamanders: genetics of ice ages
- Ensatina* 264
- E. eschscholtzii*: molecular data 273 App.
- Plethodon cinereus* 260
- Salmo*: genetics of ice ages 262, 264
- S. salar* (salmon) molecular data 275 App.
- S. trutta* (brown trout) molecular data 275 App.
- Salticidae (jumping spiders): simple eyes 39
- samples
- evenness in moth populations 75–81 & Fig. 3 & Tab. 1
- geographic cohesiveness in an intertidal snail 57, 59–62 & Fig. 1, 68–69 Figs 4–5
- SE Asian shrews 199–201 & Fig. 1
- strategy: RAPD variation in *Iliamna* 360
- Sanctacaris*: arthropod evolution 9, 24, 26, 30–31 & Fig. 2, 42
- computer-generated affinities 54–55
- sandpipers: aggressive nest defence 427
- curlew (*Calidris ferruginea*) 426, 439 App.
- upland (*Bartramia longicauda*) 426, 437 App.
- Sapindaceae: visits by Ithomiinae 109–110 & Tab. 5
- see also* *Kohliuretia*
- Sarotrocentrus*: Cambrian arthropod 14, 16, 23, 27, 37
- Savu (Indonesian archipelago): Eupreptantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
- Scandinavia: genetics of ice ages 249–250, 259
- Scaracarida: eyelessness 41 & Fig. 4
- Scatophaga stercoraria* (flies): ornaments are armaments 390
- scattergrams: flower-fruit size allometry 404–406 & Fig. 2
- Scearctia figulina* (Arctiidae: moths): pyrrolizidine alkaloids 114
- Sceloporus occidentalis* (lizards): malarial parasitic stress 474
- Sciomyzidae (Diptera): development of aquatic insect eggs 221, 231, 242 App. 2
- see also* *Ilione*, *Limnia* and *Pherbellia*
- Scolapacidae: aggressive nest defence 427
- see also* curlews, dowitchers, godwits, phalaropes, pratincoles, sandpipers, snipes, turnstones and woodcocks
- Scolopax*: aggressive nest defence
- S. minor* 438 App.
- S. rusticola* 438 App.
- scoring: evenness in moth populations 77
- Scutigera: compound eyes 38
- Scutigera: compound eyes 41 Fig. 4, 43
- sea levels: SE Asian shrews 197–198, 200 Fig. 1, 202–203, 211, 216
- seals, elephant: ornaments are armaments 389
- seedsnipe: aggressive nest defence 427
- *selection: geographic cohesiveness in an intertidal snail 57–59
- *selective sweep: mtDNA variability and male-lethal symbionts 453–455, 460, 464, 465, 467–468
- Senecio* (Asteraceae): pyrrolizidine alkaloids 100
- senecionine: (pyrrolizidine alkaloid): ithomiine butterflies 111
- sequence comparisons: arthropod evolution 1, 45–47
- Sequential Break (SB): evenness in moth populations 75, 77–78 & Fig. 1, 81
- Seram (Indonesian archipelago): Eupreptantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
- Serra do Japi, (San Paulo, Brazil): pyrrolizidine alkaloids in ithomiine butterflies 101, 106, 108–110 & Tabs 3 & 5, 112–114 & Tab. 9
- *sexual selection: conservation strategies, stress and diversity 471, 473–475
- *sexual selection: ornaments are armaments 385–392 & Tab. 1
- *sexual size dimorphism: monandry in damselflies 85, 88–90 & Fig. 1, 96
- Shannon's Index H: evenness in moth populations 76–80 & Fig. 3
- sheathbills: aggressive nest defence 427
- sheep
- bighorn (*Ovis canadensis*): horn volume 475
- mountain (*Ovis canadensis*): visual signals 391–392 Tab. 1
- Nearctic wild (*Ovis dalii*): genetics of ice ages 260
- Soay (Scotland): winter starvation 174
- shells: *Bembicium vittatum* (snails) 57, 61, 63, 70
- *shorebirds: aggressive nest protection 409–414
- shrews (*Crociodura*): Old World 199, 203, 213
- SE Asia 197–219 & Figs 1–6 & Tabs 1–4 & App.
- shrimps: sponge-dwelling alpheid 307–324 & Figs 1–5 & Tabs 1–2
- Siberia: genetics of ice ages 251–252, 258
- *sibling species: sponge-dwelling alpheid shrimps 307–308, 314
- Sibogamermes weberi* (outgroup): EIO 331
- Sicily (Mediterranean, ME): Eupreptantia 329
- Sidneyia*: extinct arthropod 16, 24, 26
- Sierra Valley; California: egg crypsis in blackbirds 164, 168
- signals: ornaments are armaments 385–393 & Tab. 1
- acoustic and visual 390–392 & Tab. 1
- *significance tests: reliability of synapomorphic analyses 483–497 & Figs 1–2 & Tabs 1–3 & Apps 1–2; Tabs A1–A4
- Silurian: arthropod evolution 9, 30, 32
- Simpson's Index: measure of diversity 79–80
- simulations: statistics of synapomorphies 487, 492–493, 496–498 App. 2 & Tab. A4
- Simulium* (blackflies): molecular data
- S. venustum* 262, 274 App.
- S. verecundum* 274 App.
- S. vittatum* 274 App.
- Siphonuridae (Ephemeroptera): development of aquatic insect eggs 241 App. 2
- see also* *Isorychia*
- Siphonaria kurracheensis* (pulmonate limpets): intertidal communities 61, 64 Tab. 1
- Siphonoperla*: (Chloroperlidae): development of aquatic insect eggs
- S. burmeisteri* 240 App. 2

- S. torrentium* 240 lp. 2
- Sirius Passet fauna: Greenland 15
- sites: geographic cohesiveness in an intertidal snail 57–70 & Figs 1 & 3 & Tab. 2 & 4, 72
- sizes
- flower-fruit allometry: *Crepis* (Asteraceae) 401–407 & Figs 1–2
 - growth similarities in tits 343–346, 352
- skuas: nest predators of waders 416, 422, 436–439 App.
- Skwala parallela* (Perlodidae): development of aquatic insect eggs 239 App. 2
- Slatkin, M. *Exploring Evolutionary Biology*. Readings from *American Scientist*. (Book Review: R. J. Berry) 500
- smelt *see Osmerus*: genetics of ice ages
- snails (*Bembicium vittatum*): geographic cohesiveness 57–74 & Figs 1–5 & Tabs 1–4
see also Austrocochlea, Bembicium, Littoraria, Litorina, Mytilus, Nerita
- snakes
- grass (*Natrix*): hybrid zones 257
- mountain garter (*Thamnophis elegans*): blackbird egg predation 162
- snipe
- aggressive nest defence
 - common (*Gallinago gallinago*) 417 Fig. 2, 438 App. seedsnipe 427
 - great (*Gallinago media*: visual signals 391–392 Tab. 1
- SOD-1, SOD-2 and SOD-3 (EC 1.15.1.1.)
- SE Asian shrews 200
- Sod-1* 204–206 Tabs 1–3
 - SOD-1 and SOD-3 203
 - SOD-2 204 Tab. 1, 206 Tab. 3
 - Sod-2* 205 Tab.
- Jalmenus evagoras*
- Sod* 299 Tab. 5
- Sod* 291, 295 Tabs 2–3, 297
- software: multivariate genetic differentiation 192
- see also ADE*
- Solanaceae: pyrrolizidine alkaloids in ithomiine butterflies 99, 101, 108–109 Tab. 4 & Fig. 3, 118–119
see also Brugmansia, Cestrum and Solanum
- Solanum*: pyrrolizidine alkaloids in ithomiine butterflies
- S. paniculatum* 101, 112
 - S. spp.* 112
- South America: genetics of ice ages 249, 257, 274 App.
- *south Devon coast: multivariate genetic differentiation 173, 175, 180–182 & Fig. 2, 187
- Southeast Asia (SE Asia): shrews (*Crocodyra*) 197, 199–201 & Fig. 1, 206–209 & Tab. 4, 213
- Spain: genetics of ice ages 250–251, 258–259, 261, 275 App.
- Spalax ehrenbergi* (mole rats)
- aridity stress 474–476
 - molecular data 275 App.
- sparrows
- genetics of ice ages
 - Melospiza* 264
 - 'seaside' (*Ammodramus maritimus*): molecular data 272 App.
 - growth: fox (*Passerella*) 352
 - ornaments are armaments
 - grasshopper (*Ammodramus swannarum*) 393
 - Harris' 387
 - house (*Passer domesticus*) 387, 391–392 Tab. 1
 - Scott's seaside (*Ammodramus maritimus*) 391–392 Tab. 1
 - white-crowned 387
 - stress: song (*Melospiza melodia*) 473
- Spearman Rank Correlation tests: aggressive nest protection by waders 420–423
- specialization: sponge-dwelling alpheid shrimps 307–309, 321
- speciation: genetics of ice ages 247–248, 254–257, 264, 266
- *speciation: sponge-dwelling alpheid shrimps 307–308, 321
- Speciation and the Recognition Concept, Theory and Application*, by D. M. Lambert and H. G. Spencer. (M. F. Claridge. Book Review) 499
- species
- boundaries: specialization and speciation in shrimps 307–309, 312–316 & Fig. 1, 319–321
 - development of aquatic insect eggs 221–223, 225–228, 230–233, 238–242 App. 2
 - first fossil anthomyid 441, 447–450
 - genetics of ice ages 247–253 & Fig. 1, 256–266 & Fig. 7, 272–276 App.
 - growth similarities in tits 343–347, 349–354 & Tab. 4
 - mtDNA variability and male-lethal symbionts 453–454, 457
 - RAPD variation in *Iliamna* 357–360, 364–365
 - see also* endangered species
- *species richness: complementary areas for diversity 125–138 & Figs 1 & 3 & 6 & Tabs 1–2, 144–146 App. 1, 152–154 App. 3
- spectrometry: pyrrolizidine alkaloids (PAs) in ithomiine butterflies 99, 105
- *sperm competition: monandry in damselflies 85–87, 95–97
- Sphingidae (hawkmoths): colour polymorphism 278
- spiders
- arthropod evolution
 - camera-type eyes 37, 42
 - jumping (Salticidae): simple eyes 39
 - labral development 16
 - molecular data: *Tetragnatha* 273 App.
 - ornaments are armaments
 - Linyphia litigiosa* 390
 - social (*Acheareanea wau*) 391–392 Tab. 1
 - wandering (*Cupiennius gelazi*) 391–392 Tab. 1
- pyrrolizidine alkaloids (PAs) 120
- orb weaving (*Nephila clavipes*): predation of ithomiine butterflies 99–101, 115, 118–121
- spines (on penes)
- monandry in damselflies 85, 88, 91 Tab. 1, 95, 97
 - Sympetrum danae* (anisopteran) 95
- Spodoptera littoralis*: mtDNA variability and male-lethal symbionts 467
- sponges: specialization and speciation in shrimps 307–318 & Figs 1 & 3, 321
see also Agelas, Hyatella, Hymeniacidon, Lissodendoryx, Oceanapia and Xestospongia
- spruce budworm: mtDNA divergence 266
- squirrels, California ground (*Citellus beecheyi*): blackbird egg predation 162

- Stachytarpheta cayennensis* (Verbenaceae): *Aeria olena* visits 106–108 Fig. 2 & Tabs 1 & 3
- starlings, European (*Sturnus vulgaris*): effects of competition 478
- Start Bay/Point (Devon coast): *Nucella lapillus* (dogwhelks) 181–190 & Figs 2–3 & 5
- Statistica 4.0: aggressive nest protection by waders 423
- statistical analyses
- cladistic biogeography of the Eureptantia 337 Tab. 3, 339
 - see also analyses and computer programs
 - development of aquatic insect eggs 224, 236–242
 - Apps 1–2
 - see also analyses and regressions
 - effect of light and diet on a polymorphic caterpillar 280–281
 - see also model I ANOVA
 - evenness in moth populations 75–81 & Figs 1–2 & Tab. 1
 - see also indices, log distributions and indices
 - flower-fruit size allometry 405
 - see also correlations, major axis regression and null hypothesis
 - geographic cohesiveness in an intertidal snail 75–80 & Figs 1–3 & Tabs 1–4
 - see also Bonferroni, chi-squares, computer programs, correlations, indices, Nei and regressions
 - monandry in damselflies 86–87, 89
 - see also Kruskal-Wallis, principal components and regressions
 - multivariate genetic differentiation 174, 178 179 & Tab. 1, 187, 194–195 App.
 - see also analyses, eigenvalues, computer programs, correlations, Hardy-Weinberg and Student's *t*-tests
 - ornaments are armaments 392
 - RAPD variation in *Iliamna* 357, 359–361, 364–365 & Fig. 3 & Tab. 3
 - see also AMOVA, F-statistic, HOMOVA, Mantel test and UPGMA
 - *statistics: reliability of synapomorphic analyses 483–497 & Figs 1–2 & Tabs 1–3 & Apps 1 2:- Tabs A1–A4
 - basic assumptions 186 187
 - see also arcsine transformations, covariances and Freeman-Tukey method
 - status: ornaments are armaments 385–390
 - Statview SE + Graphics: geographic cohesiveness in an intertidal snail 61–62
 - Stephanopodium*: complementary areas for diversity
 - S. aptotum* 151 App. 2
 - S. engleri* 150 App. 2
 - S. gentryi* 149 App. 2
 - S. magnifolium* 147 App. 2
 - S. sessile* 147 App. 2
 - S. venezuelanum* 151 App. 2 - sticklebacks (*Gasterosteus*)
 - genetics of ice ages 264
 - G. aculeatus*: molecular data 272 App.
 - ornaments are armaments: three-spined (*G. aculeatus*) 388, 391–392 Tab. 1 - stilt: aggressive nest defence 422
 - black-winged (*Himantopus himantopus*) 417 Fig. 2, 436 App.
 - stingrays: helminth parasites 340
 - stoneflies see Plecoptera
 - strategies: conservation, stress and diversity 471–472, 477–478
 - stress: conservation strategies and diversity 471–460
 - striders, water: cladistic biogeography 340
 - Student's *t*-test: multivariate genetic differentiation 189–190 & Fig. 7
 - study areas
 - egg crypsis in blackbirds 164
 - pyrrolizidine alkaloids in ithomiine butterflies 101
 - see also Amarais wood, Costa e Silva wood, Mata da Pedra, Monjolinho Arboretum and Serra do Japi
 - sturgeon, white (*Acipenser transmontanus*): molecular data 272 App.
 - Sturnus vulgaris* (European starlings): effects of competition 478
 - subpopulations: butterfly-ant mutualism 287
 - subspecies: RAPD variation in *Iliamna* 357
 - Sulawesi
 - Eastern Indian Ocean (EIO): Eureptantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4 - SE Asian shrews 197, 199–200 & Fig. 1, 202–203 & Fig. 2, 206–209 & Tab. 3, 211–216 & Figs 5–6, 219 App.
 - New Sulawesi 210–212 & Fig. 6
 - Sumatra: SE Asian shrews 197–198, 200 Fig. 1, 202–203 & Fig. 2, 205 Tab. 2, 207–212 & Figs 5–6, 214–215, 219 App.
 - Sumba (Indonesian archipelago): Eureptantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
 - summary: cladistic biogeography of the Eureptantia 340–341
 - Suncus dayi* (shrews): India 199, 201, 203–204 & Tab. 1, 208–211 Tab. 4 & Figs 3–4
 - Sunda Shelf: SE Asian shrews 197–199, 202 & Fig. 2, 207–210, 211–215 & Figs 5–6
 - sunfish (*Lepomis*): genetics of ice ages 264
 - bluegill (*L. macrochirus*): molecular data 272 App.
 - rederar (*L. microlophus*) 272 App.
 - spotted (*L. punctatus*) 272 App.
 - Warmouth (*L. gulosus*) 272 App.
 - supinine (pyrrolizidine alkaloid): ithomiine butterflies 99, 102 Fig. 1, 104, 114
 - survival
 - egg crypsis in blackbirds 159, 162, 167 168 & Tab. 3 & Fig. 4, 170–171
 - rates (damselflies)
 - Fisher and Ford's Method 92–93, 95
 - Jolly-Seber Stochastic Method 92
 - Manly-Parr estimates 92–93 - suture zones: genetics of ice ages 247, 257–258, 262
 - Sweden (Uppsala): growth similarities in tits 345–346
 - swordtails, green (*Xiphophorus helleri*): visual signals 391–392 Tab. 1
 - symbionts, male-lethal: mtDNA variability 453–470 & Figs 1–6 & Tabs 1–3 & App
 - *symbiosis: specialization and speciation in shrimps 307, 321
 - Sympetrum* (anisopteran)
 - development of aquatic insect eggs 233
 - paired spines on accessory penes: *S. danae* 95 - Symphyla: eyeless or without compound eyes 41 Fig. 4

- **Synalpheus*: sponge-dwelling alpheid shrimps 307–309, 314, 320–321
S. brooksi 310, 320–321
S. fidalgus 307–308, 314–316, 318–319 & Fig. 4, 321
S. rathbunae 307–309, 312–322 & Figs 1–5 & Tabs 1–2
 A 321
 A, B and C 312–320 & Figs 1–4 & Tab. 2
 *synapomorphies: statistical reliability of analyses 483–497 & Figs 1–2 & Tabs 1–3 & Apps 1–2 including Tabs A1–A4
 Systat version 5.2: geographic cohesiveness in an intertidal snail 63
 Systelognatha (Plecoptera): development of aquatic insect eggs 221, 230, 239–240 App. 2
see also Chloroperlidae, Perlidae and Perlodidae
 systematics: first fossil anthomyid 442–447 & Figs 1–2
- Taeniopterygidae (Plecoptera): development of aquatic insect eggs 239 App. 2
Taeniopteryx nebulosa 224 Fig. 1, 239 App. 2
 tandem guarding: monandry in damselflies 85–88, 90–91 & Tab. 1, 95
 Tantulacarida: eyeless or without compound eyes 41–42 & Fig. 4
Tapura: complementary areas for diversity
T. bullata 149 App. 2
T. costata 148 App. 2
T. cubensis s. minor 151 App. 2
T. ferreyrae 149 App. 2
T. haihensis 151 App. 2
T. lanceolata 148 App. 2
T. magnifoia 150 App. 2
T. mexicana 149 App. 2
T. orbicularis 151 App. 2
T. panamensis 150 App. 2
T. tessmannii 151 App. 2
 Tardigrada: arthropod evolution 12, 15, 41 Fig. 4
 *taxon: first fossil anthomyid and comments on clade definitions 441, 447–449
- taxonomy
 aggressive nest protection by waders 417–418, 436–439 App
 Eureptantia 326–327 & Fig. 1
Iliamna corei and *I. remota* 358–361
 SE Asian shrews 213
Tellera zolius: ithomiine butterflies 118
 temperatures: development of aquatic insect eggs 221–233 & Figs 1–2 & 4–5, 236–237 App. 1
 temporal changes: evenness in moth populations 78–79 & Fig. 2
 Tennessee: beech pollen cores 250–251
 tension zones: genetics of ice ages 256
 terrapins, diamondback (*Malaclemys terrapin*): molecular data 272 App.
 Tertiary
 genetics of ice ages 249
 Indian collision with the Asian continent: Eureptantia 340
 Tethys Sea: cladistic biogeography of the Eureptantia 325, 338–340
Tetragnatha (spiders): molecular data 273 App.
Tetraopes tetraophthalmus (cerambycid beetle): extinction/recolonization 303
 tetrapods: statistics of synapomorphies 483, 489–491 & Tabs 4–5
 Texas: *Ischnura hastatum* and *I. kellicotti* 87
 thaid *Nucella lapillus*: localised selection 58
Thalestria rhodymeniae: arthropod evolution 13
Thamnophis elegans (mountain garter snakes): blackbird egg predation 162
 Theria: statistics of synapomorphies 490–492 & Tabs 2–3, 496 Apps 1–2 & Tab. A4
 *thermal reaction norms: development of aquatic insect eggs 221–233 & Figs 1–5
 thick-knees: aggressive nest defence 427
see also *Burhinus*
 Thomason, J. J. (ed.). *Functional Morphology in Vertebrate Palaeontology* (Book Review: Peter Forey) 245–246
Thomomys (pocket gophers): genetics of ice ages 259, 264
T. bottae 263, 274 App.
T. umbrinus 263, 274 App.
 three area statements: cladistic biogeography of the Eureptantia 326, 336–337 & Figs 5–6
Thylacinus (marsupial): convergence 35
 Thysanura: arthropod evolution 9, 41 Fig. 4, 43
 Timor (Indonesian archipelago): Eureptantia 327–328 & Fig. 1, 331, 333 Tab. 2, 335 Fig. 4
 Tipulidae (Diptera): development of aquatic insect eggs 221, 231, 242 App. 2
Tipula 231
T. commiscibilis 242 App. 2
T. maxima 242 App. 2
 Tipulinae 231
 **Tithorea harmoniaplain* (ithomiine butterflies): pyrrolizidine alkaloids 99, 101, 104, 108–111 & Tabs 4 & 6 & Fig. 3, 113–115, 118–119
T. h. pseudethra 101
- tits
 great (*Parus major*): ornaments are armaments 387, 391 392 Tab. 1
 growth similarities 343–345, 351, 353–354
see also blue and great tits
 Tiur (Indonesian archipelago): Eureptantia 327–328 & Fig. 1, 333 Tab. 2, 335 Fig. 4
 toadfish (*Opsanus*): genetics of ice ages 264
O. beta: molecular data 272 App.
 toads
 natterjack (*Bufo calamita*): acoustic signals 391–392 Tab. 1
see also *Bombina*: hybrid zones
 tortoises, gopher (*Xerobates*): molecular data
X. agassizi 273 App.
X. berlandieri 273 App.
Toxorhynchites (Culicidae): development of aquatic insect eggs
T. brevipalpis 224 Fig. 1, 242 App. 2
T. rutilis 242 App. 2
Tpi (enzyme loci): sponge-dwelling alpheid shrimps 310
 Tracheata: arthropod evolution 1–2, 6, 42
 early: comparison with ancestral crustaceans 20–22
see also Hexapoda and Myriapoda
 traits
 first fossil anthomyid 441, 445–447
 flower-fruit size allometry 401–402
 growth similarities in tits 343, 345–353 & Tabs 1–6 & Fig. 1

- ornaments are armaments 385–394 & Tab. 1
- transmission efficiency: mtDNA variability and male-lethal symbionts 455, 451–462, 464–466
- Transylvania: genetics of ice ages 249
- treehoppers (*Enchenopa binotata*): genetics of ice ages 266
- trees: genetics of ice ages 250–251, 257
 see also alder, beech, birch and hazel
- triangularine: (pyrrolizidine alkaloid): ithomiine butterflies 114
- Trichogonia gardneri* (Asteraceae: Eupatoriaceae): pyrrolizidine alkaloids in ithomiine butterflies 106, 108 Tab. 3, 110 Tab. 5, 113 & Tab. 9, 117
- Trichopsis vittatus* (croaking gura): acoustic signals 391–392 Tab. 1
- Tricorythidae (Ephemoptera): development of aquatic insect eggs 242 App. 2
- Tricorythodes minutus* 242 App. 2
- Trilobita: arthropod evolution 2, 4–6, 14, 23, 41 Fig. 4
 polyphyletic concept 25 Fig. 1, 31 Fig. 2
- Tringa*: aggressive nest protection 418
T. erythropus 418, 437 App.
T. flavipes 427, 438 App.
T. glariola 438 App.
T. guttifer 437 App.
T. melanoleuca 437 App.
T. nebularia 437 App.
T. ochropus 438 App.
T. solitaria 438 App.
T. stagnatilis 437 App.
T. totanus (redshank) 426, 437 App.
- Trinidad: *Poecilia reticulata* (guppies) 272 App.
- Triops*: arthropod evolution 3
- Triturus* (newts)
 hybrid zone barriers 259, 261
T. carnifex: molecular data 275 App.
T. cristatus (crested newts) 260–261, 275 App.
T. dobrogicus 275 App.
T. karelini 275 App.
T. marmoratus 275 App.
- trochid *Austrocochlea constricta*: geographic cohesiveness in an intertidal snail 61, 63
- trout
 rainbow (*Onchorhynchus mykiss*): bacterial gill disease 474
Salmo: genetics of ice ages 264
 brown (*S. trutta*) 275 App.
- Tryngites subruficollis*: aggressive nest defence 439 App.
- Tukey's multiple comparisons: development of aquatic insect eggs 228
- Turkey: genetics of ice ages 251–252, 258–259, 261, 264
- turnstones: aggressive nest protection 422, 427
 see also *Arenaria*
- twigs, oak: effect of light and diet on a polymorphic caterpillar 277, 279–283 & Tab. 1
- Uca pugilator* (fiddler crabs): ornaments are armaments 386, 391–392 Tab. 1
- umbellifer hosts: *Greya suffusa* (moths) 266
- ungulates: ornaments are armaments 386
 see also antelopes, bison, caribou, deer and pronghorns
- Uniporus*: eurentantic nemerteans
- U. acutocaudatus* (EAT) 327 Fig. 1, 329, 332 Fig. 2
- U. borealis* (EAT) 327 Fig. 1, 329, 332 Fig. 2
- U. hyalinus* (EAT) 327 Fig. 1, 329, 332 Fig. 2
- Uniramia: arthropod evolution 2, 5, 25 Fig. 1, 31 Fig. 2, 41 Fig. 4
 Uniramia Vera 20, 42, 49
 validity of the taxon 45, 49
- Uperoleia rugosa* (Australian frogs): acoustic signals 391–392 Tab. 1
- UPGMA (unweighted paired group method using arithmetic averages)
 RAPD variation in *Iliamna* 357, 361, 365
 SE Asian shrews 201–203 & Fig. 2, 208–209
 specialization and speciation in shrimps 310, 312 Fig. 1
- Uppsala, Sweden: growth similarities in tits 345–346
- Ur-Crustacea: arthropod evolution 19–21, 40–41 & Fig. 4
- Uralia*: arthropod evolution 5, 9–10, 16
- Urals: genetics of ice ages 257
- Urbanus proteus* (Hesperiidae): non-pyrrolizidine alkaloid insects 109
- urchins (*Echinometa mathaei*): intertidal communities 61, 64 Tab. 1
- Urichonemertes pilorhynchus*: eurentantic nemertean (WP: Gbr) 327–329 & Fig. 1, 332 Fig. 2, 339
- USA
 genetics of ice ages 262–263, 272–274 App.
Quercus (oaks) 178
 see also Arizona, Florida, Illinois, Nevada, New Mexico, Texas, Tennessee and Virginia
- usaramine (pyrrolizidine alkaloid): ithomiine butterflies 111
- Utacapnia trava* (Capniidae): development of aquatic insect eggs 238 App. 2
- Utiethesia ornatrix* (Arctiidae: moths): chemical defences 118
- Vaccinium macrocarpon* (Ericaceae): use of AMOVA 360
- vallylleucine peptidase (EC 3.4.1.3: *Bembicium vittatum* (snails)
Vlp1 65
Vlp1 and *Vlp2* loci) 62, 65, 72
Vlp2 65, 67
- Vanellus*: aggressive nest protection
V. cinereus 436 App.
V. gregarius 436 App.
V. indicus 436 App.
V. lecurus 436 App.
V. malabaricus 413
V. sponus 436 App.
V. vanellus (lapwings) 411, 417 Fig. 2, 422, 436 App.
- variances
 growth similarities in tits 346–351 & Tabs 1–2 & 5
 RAPD variation in *Iliamna* 357, 359–360, 364–365 & Fig. 2
 see also ANOVA and HOMOVA
- variations
 aggressive nest protection by waders 409–411, 415, 418, 423–428
 mtDNA and male-lethal symbionts 453–455
- vein CuP ('anal vein'): diagnosis of first fossil anthomyid 442, 445–448

- Vendian times; late Pre-Cambrian arthropod evolution 26
- Verbenaceae: *Aeria olen* visits 106–108 Fig. 2 & Tabs 1 & 3
see also Lantana and *Stachytarpheta*
- Vernoniaceae (Asteraceae): pyrrolizidine alkaloids in ithomiine butterflies 106 Fig. 2, 109
Vernonia polyanthes 106 Fig. 2, 110 Tab. 5
V. rubriramea 109, 110 Tab. 5
see also Elephantopus
- viability: complementary areas for diversity 126, 139–140
- Violaceae: *Aeria olen* visits 106–107 Fig. 2 & Tab. 1
see also Hybanthus and *Jonidium*
- Virginia: RAPD variation in *Iliamna* 357–363 & Fig. 1 & Tabs 1–2, 365
see also Blackburn, Peters Mountain, populations (*Iliamna*)
- Vulpes* (foxes)
 genetics of ice ages 261
 molecular data
V. macrotis (kit fox) 274 App.
V. velox (swift fox) 274 App.
- waders, Holarctic: aggressive nest defence 409–429 & Figs 4–6 & Tabs 1–2, 436–439 App.
- Wagner trees: SE Asian shrews 201, 207, 210 Fig. 3
 distance-Wagner 201, 206
- Wallabi Group (Houtman Albrohous Islands, Western Australia): *Bembicium vittatum* (snails) 60 Fig. 1, 63, 65 & Fig. 2, 67–70 & Tabs 2 & 4 & Fig. 4, 72
- warblers, great reed (*Acrocephalus arundinaceus*): acoustic signals 391–393 & Tab. 1
- wasps, oak gall (*Andricus quercuscalicis*): genetics of ice ages 260
- water depths: SE Asian shrews 197–198, 200 Fig. 1, 202–203 & Fig. 2, 207–209, 215–216
- *weapon badge: ornaments are armaments 385–386, 394
see also armaments
- weavers, village: egg coloration 169
- whimbrels (*Numenius phaeopus*): aggressive nest protection 411, 437 App.
- whitefish, lake (*Coregonus dupeaformis*): genetics of ice ages 261, 272 App.
- widowbirds (*Euplectes*): visual signals
 long-tailed (*E. progne*) 391–392 Tab. 1
 yellow-shouldered (*E. macrurus*) 391–392 Tab. 1
- Wilcoxon signed-rank test: egg crypsis in blackbirds 167 & Tab. 3
- Wilks' lambda: morphometric analysis of shrimp populations 313, 316 Fig. 2
- par wing lengths: monandry in damselflies 89–91 & Fig. 1 & Tab. 1
- wings: aggressive nest protection by waders
 loading 409, 411, 415–417 & Figs 1–3, 422, 424–425 & Tabs 1–2, 428
 structure 409, 416–417 & Figs 1–2, 424–425 & Tabs 1–2, 428, 436–439 App.
- Wolbachia (symbiont): mtDNA variability 453–454, 468
- wolves (*Canis lupus*): molecular data 273 App.
- Woodchester Park Field Centre; southern England: evenness in moth populations 75–76, 79 Fig. 2, 82
- woodcocks: aggressive nest defence 427
see also Scolopax
- woodland: evenness in moth populations 75–76, 80–82 & Tab. 1
- woodpeckers (*Colaptes auratus*): genetics of ice ages 263
- woodrats, eastern (*Neotoma*): genetics of ice ages 264
N. albigula: molecular data 274 App.
N. floridana 261, 264, 274 App.
N. micropus 274 App.
- Workshop 90 (Biological Priorities for Conservation in Amazonia): complementary areas for diversity 125–130, 134–139 & Fig. 1 & Tabs 1–2
- WORLDMAP software: complementary areas for diversity 130, 138–140
- worms
 arthropod evolution
 flatworms 25 Fig. 1
 worm-like ancestors 10, 12, 14, 17, 26, 28
 heterozygote advantage: manure worms (*Eisenia foetida*) 474
- *Wright's F-statistics: population genetic effects of a butterfly-ant mutualism 287, 292–296 & Fig. 2 & Tabs 2–4, 298–299, 302
- Wulffia* spp. (Asteraceae): *Aeria olen* visits 108 Tab. 3
- Xanthocephalus xanthocephalus* (yellow-headed blackbirds)
 egg crypsis 159–162 & Fig. 1, 164–171 & Tab. 1–3 & Figs 2–4
 visual signals 391–392 Tab. 1
- XDH: SE Asian shrews 200–201, 203
- Xenonemertes rhamphocephalus*: eurentatic nemertean (WP: Gbr) 327–329 & Fig. 1, 332 Fig. 2, 339
- Xenus cinereus*: aggressive nest defence 438 App.
- Xenusion*: Cambrian fossils 15
- Xerobates* (gopher tortoises): molecular data
X. agassizi 273 App.
X. berlandieri 273 App.
- Xestospongia* (sponges) 321
X. cf. proxima 316, 318 Fig. 4
X. cf. subtriangularis: Belize 309–310, 312–318 & Figs 1 & 4
X. rosariensis Panama 309–315 & Fig. 1, 318–319 & Figs 4–5
- Xiphophorus helleri* (green swordtails): visual signals 391–392 Tab. 1
- Xiphosura: arthropod evolution 30
- yellowlegs, lesser (*Tringa flavipes*): aggressive nest defence 427
- Younger Dryas: genetics of ice ages 250–251
- Zahavi's handicap principle: ornaments are armaments 388
- Zapada* (Nemouridae): development of aquatic insect eggs
Z. cinctipes 239 App. 2
Z. columbiana 239 App. 2
- Zea mays*: inbreeding depression 473
- *zoogeography: SE Asian shrews 197, 202–203 & Fig. 2, 214–216
- Zygoptera: monandry in damselflies 86

